

CUT TIME IN SUB-RIEMANNIAN PROBLEM ON ENGEL GROUP

A. A. ARDENTOV¹ AND YU. L. SACHKOV¹

Abstract. The left-invariant sub-Riemannian problem on the Engel group is considered. The problem gives the nilpotent approximation to generic nonholonomic systems in four-dimensional space with two-dimensional control, for instance to a system which describes motion of mobile robot with a trailer.

The global optimality of extremal trajectories is studied via geometric control theory. The global diffeomorphic structure of the exponential mapping is described. As a consequence, the cut time is proved to be equal to the first Maxwell time corresponding to discrete symmetries of the exponential mapping.

Résumé. ...

1991 Mathematics Subject Classification. 22E25, 58E25.

28.08.2014.

INTRODUCTION

This paper continues the study of the left-invariant sub-Riemannian problem on the Engel group started in [1, 2]. This problem is the simplest rank 2 sub-Riemannian problem on a 4-dimensional space: it provides a nilpotent approximation to a generic sub-Riemannian problem of such kind near a generic point.

A sub-Riemannian (SR) structure on a smooth manifold M is a vector distribution

$$\Delta = \{\Delta_q \subset T_q M \mid q \in M\} \subset TM$$

with a scalar product in Δ :

$$g = \{g_q \text{ — scalar product in } \Delta_q \mid q \in M\}.$$

The subspaces $\Delta_q \subset T_q M$ and the scalar product $g_q: \Delta_q \times \Delta_q \rightarrow \mathbb{R}$ depend smoothly on a point $q \in M$. The dimension of the subspaces Δ_q is constant ($\dim \Delta_q$ is called the rank of the distribution Δ).

A Lipschitz curve $q: [0, t_1] \rightarrow M$ is horizontal if $\dot{q}(t) \in \Delta_{q(t)}$ for almost all $t \in [0, t_1]$. The length of a horizontal curve is $l = \int_0^{t_1} g(\dot{q}(t), \dot{q}(t))^{1/2} dt$. The sub-Riemannian distance between points $q_0, q_1 \in M$ is the infimum of lengths of horizontal curves that connect q_0 to q_1 . A horizontal curve $q(t), t \in [0, t_1]$, is a (length) minimizer if it has a minimum possible length among all horizontal curves that connect the points $q(0)$ and

Keywords and phrases: sub-Riemannian geometry, optimal control, Engel group, Lie algebra, Maxwell time, cut time, exponential mapping, Euler's elastica

¹ Program Systems Institute of RAS, Pereslavl-Zalessky, Russia; e-mail: aaa@pereslavl.ru yusachkov@gmail.com

$q(t_1)$. Description of minimizers is one of important problems of sub-Riemannian geometry. The most efficient approach to this problem is given by geometric control theory [3–5], it consists of the following steps:

- (1) proof of existence of minimizers,
- (2) description of SR geodesics (*i.e.*, curves whose small arcs are minimizers),
- (3) selection of minimizers among geodesics.

Step (1) is straightforward. If M is connected and Δ is bracket generating, *i.e.*, $\text{Lie}_q \Delta = T_q M$, $\forall q \in M$, then any points $q_0, q_1 \in M$ can be connected one to another by a horizontal curve (Rashevsky-Chow theorem). If additionally the point q_1 is sufficiently close to q_0 , or if the SR distance is complete, or if Δ and g are left-invariant on a Lie group M , then q_0 can be connected with q_1 by a minimizer (Filippov theorem).

Step (2) is performed via application of Pontryagin maximum principle (PMP), which states that any geodesic (thus any minimizer) is a projection of a trajectory of a certain Hamiltonian system on the cotangent bundle T^*M . So the second step reduces to the study of integrability of the Hamiltonian system of PMP and efficient parameterization of trajectories of this system.

Step (3) is the hardest one. Local optimality of geodesics (*i.e.*, optimality w.r.t. sufficiently close geodesics) is studied via conjugate points estimates. For the study of global optimality in problems with a big symmetry group, one can often obtain bounds (or explicit description) of cut time via the study of symmetries and global structure of the exponential mapping. We suggest the following detailing of Step (3) first applied in [6] and further developed in [7–16]:

- (3.1) Discrete and continuous symmetries of the exponential mapping are found;
- (3.2) Maxwell points corresponding to the symmetries are found (*i.e.*, points where several geodesics obtained one from another by a symmetry meet one another). These points (and their preimage via exponential mapping) form the Maxwell strata in the image (resp., in the preimage) of the exponential mapping. Along each geodesic, the first Maxwell time corresponding to the symmetries (*i.e.*, the time when the geodesic meets a Maxwell strata) is found;
- (3.3) One proves that for any geodesic the first conjugate time is greater or equal to the first Maxwell time corresponding to the symmetries. Here the homotopy invariance of Maslov index (number of conjugate points on a geodesic) can be applied [17];
- (3.4) One considers restriction of the exponential mapping to the subdomains cut out in preimage and image of this mapping by the Maxwell strata corresponding to symmetries, and proves that this restriction is a diffeomorphism via Hadamard global diffeomorphism theorem [18];
- (3.5) On the basis of the global structure of the exponential mapping thus described, it is often possible to prove that the cut time along a geodesic (*i.e.*, time when it loses its global optimality) is equal to the first Maxwell time corresponding to symmetries. Moreover, in this way one proves that for any terminal point in a subdomain in the image of the exponential mapping, there exists a unique minimizer which can be computed by inverting the exponential mapping in the subdomain;
- (3.6) Finally, for systems with big symmetry group one can construct the full optimal synthesis, and numerical algorithms and software for computation of optimal trajectories with given boundary conditions.

So far, the approach described has been applied in full just to several problems: SR problem in the flat Martinet case [6], SR problems on $\text{SO}(3)$ and $\text{SL}(2)$ with the Killing metric [19], SR problem on $\text{SE}(2)$ [11–13], Euler elastic problem [14–16]. There are partial results on the nilpotent SR problem with the growth vector $(2, 3, 5)$ [7–10] and SR problem on $\text{SH}(2)$ [20, 21].

For the SR problem on the Engel group, Step (1), Step (2) and Steps (3.1), (3.2) are performed in [1] while (3.3) is done in [2]. The aim of this paper is to perform Steps (3.4), (3.5). We recall the results previously obtained in the next section.

The sub-Riemannian problem on the Engel group is a left-invariant problem on a Lie group. Such problems receive significant attention in geometric control since they provide very symmetric models which can often be studied explicitly in great detail. For left-invariant SR problems on Lie groups, one can often describe optimal synthesis, the structure of spheres, cut and conjugate loci. This information can give insight for general problems, where such a detailed study is much more complicated.

Left-invariant SR problems on 3D and 4D Lie groups have recently been fully classified [22, 23]. In the 3-dimensional case, optimal synthesis is known for the Heisenberg group [24], for $SO(3)$ and $SL(2)$ with the Killing metric [19] and for $SE(2)$ [11–13]. This work continues a detailed study of the simplest 4-dimensional case.

1. PREVIOUSLY OBTAINED RESULTS

In this section we recall results on the SR problem on the Engel group obtained previously in works [1, 2].


1.1. Problem statement

The Engel group is the 4-dimensional Lie group represented by matrices as follows:

$$M = \left\{ \left(\begin{array}{cccc} 1 & b & c & d \\ 0 & 1 & a & a^2/2 \\ 0 & 0 & 1 & a \\ 0 & 0 & 0 & 1 \end{array} \right) \mid a, b, c, d \in \mathbb{R} \right\}.$$

It is a 4-dimensional nilpotent Lie group, connected and simply connected (see an explanation of the name "Engel" for this group in [25], Sec. 6.11).

The Lie algebra of the Engel group is the four-dimensional nilpotent Lie algebra $L = \text{span}(X_1, X_2, X_3, X_4)$ with the multiplication table



$$\begin{aligned} X_3 &= [X_1, X_2], \\ X_4 &= [X_1, X_3], \\ [X_2, X_3] &= [X_1, X_4] = [X_2, X_4] = 0. \end{aligned} \tag{1}$$

Thus it has graduation

$$\begin{aligned} L &= L_1 \oplus L_2 \oplus L_3, \\ L_1 &= \text{span}(X_1, X_2), \quad L_2 = \mathbb{R}X_3, \quad L_3 = \mathbb{R}X_4, \\ [L_i, L_j] &= L_{i+j}, \quad L_k = \{0\} \text{ for } k \geq 4, \end{aligned}$$

and the Engel group is a Carnot group [25].

We consider the sub-Riemannian problem on the Engel group M for the left-invariant sub-Riemannian structure generated by the orthonormal frame X_1, X_2 :

$$\begin{aligned} \dot{q} &= u_1 X_1(q) + u_2 X_2(q), \quad q \in M, \quad (u_1, u_2) \in \mathbb{R}^2, \\ q(0) &= q_0, \quad q(t_1) = q_1, \\ l &= \int_0^{t_1} \sqrt{u_1^2 + u_2^2} dt \rightarrow \min. \end{aligned}$$

In appropriate coordinates $q = (x, y, z, v)$ on the Engel group $M \cong \mathbb{R}^4$, the problem is stated as follows:

$$\dot{q} = \begin{pmatrix} \dot{x} \\ \dot{y} \\ \dot{z} \\ \dot{v} \end{pmatrix} = u_1 \begin{pmatrix} 1 \\ 0 \\ -y/2 \\ 0 \end{pmatrix} + u_2 \begin{pmatrix} 0 \\ 1 \\ x/2 \\ (x^2 + y^2)/2 \end{pmatrix}, \quad q = (x, y, z, v) \in M = \mathbb{R}^4, \quad (u_1, u_2) \in \mathbb{R}^2, \quad (2)$$

$$q(0) = q_0 = (x_0, y_0, z_0, v_0), \quad q(t_1) = q_1 = (x_1, y_1, z_1, v_1), \quad (3)$$

$$l = \int_0^{t_1} \sqrt{u_1^2 + u_2^2} dt \rightarrow \min. \quad (4)$$

By virtue of the multiplication table (1) for the vector fields of the orthonormal frame

$$X_1 = \frac{\partial}{\partial x} - \frac{y}{2} \frac{\partial}{\partial z}, \quad X_2 = \frac{\partial}{\partial y} + \frac{x}{2} \frac{\partial}{\partial z} + \frac{x^2 + y^2}{2} \frac{\partial}{\partial v}$$

and their Lie brackets

$$X_3 = [X_1, X_2] = \frac{\partial}{\partial z} + x \frac{\partial}{\partial v}, \quad X_4 = [X_1, X_3] = \frac{\partial}{\partial v},$$

system (2) is completely controllable, *i.e.*, any points $q_0, q_1 \in \mathbb{R}^4$ can be connected by its trajectory.

Since the problem is invariant under left shifts on the Engel group, we can assume that the initial point is the identity $q_0 = (x_0, y_0, z_0, v_0) = (0, 0, 0, 0)$.

1.2. Parameterization of geodesics

Existence of optimal solutions of problem (2)–(4) is implied by Filippov theorem [4]. By the Cauchy-Schwarz inequality, it follows that sub-Riemannian length minimization problem (4) is equivalent to the action minimization problem:

$$\int_0^{t_1} \frac{u_1^2 + u_2^2}{2} dt \rightarrow \min. \quad (5)$$

Pontryagin maximum principle [4, 26] was applied to the resulting optimal control problem (2), (3), (5) in [1].

A sub-Riemannian geodesic can be normal or abnormal, or both. For the SR problem on the Engel group, each abnormal geodesic is simultaneously normal, thus in the sequel we consider only normal geodesics.

Normal geodesics are projections $q_t = \pi(\lambda_t)$ via the canonical projection $\pi: T^*M \rightarrow M$ of solutions to the Hamiltonian system

$$\dot{\lambda} = \vec{H}(\lambda), \quad \lambda \in T^*M, \quad (6)$$

with the Hamiltonian function $H = \frac{1}{2}(h_1^2 + h_2^2)$. Here and below $h_i(\lambda) = \langle \lambda, X_i(q) \rangle$, $\lambda \in T^*M$, $i = 1, \dots, 4$, are Hamiltonians that correspond to the left-invariant frame and are linear on fibers of the cotangent bundle T^*M .

Arclength parameterized geodesics (*i.e.*, with velocity $g(\dot{q}_t, \dot{q}_t) \equiv 1$) are projections of extremals λ_t lying on the level surface $\{\lambda \in T^*M \mid H(\lambda) = 1/2\}$.

Introduce coordinates (θ, c, α) on the level surface $\{\lambda \in T^*M \mid H = 1/2\}$ by the following formulas:

$$h_1 = \cos(\theta + \pi/2), \quad h_2 = \sin(\theta + \pi/2), \quad h_3 = c, \quad h_4 = \alpha.$$

On this surface the normal Hamiltonian system (6) takes the following form:

$$\begin{aligned} \dot{\theta} &= c, & \dot{c} &= -\alpha \sin \theta, & \dot{\alpha} &= 0, \\ \dot{q} &= \cos \theta X_1(q) + \sin \theta X_2(q), & q(0) &= q_0. \end{aligned} \quad (7)$$

The family of all normal extremals is parameterized by points of the phase cylinder of pendulum

$$C = \{\lambda \in T_{q_0}^* M \mid H(\lambda) = 1/2\} = \{(\theta, c, \alpha) \mid \theta \in S^1, c, \alpha \in \mathbb{R}\},$$

and is given by the exponential mapping

$$\begin{aligned} \text{Exp}: N = C \times \mathbb{R}_+ &\rightarrow M, \\ \text{Exp}(\lambda, t) &= q_t = (x_t, y_t, z_t, v_t). \end{aligned}$$

The energy integral of pendulum (7) is given by $E = \frac{c^2}{2} - \alpha \cos \theta$. The cylinder C has the following stratification corresponding to the particular type of trajectories of the pendulum:

$$C = \bigcup_{i=1}^7 C_i, \quad C_i \cap C_j = \emptyset, \quad i \neq j, \quad \lambda = (\theta, c, \alpha), \quad (8)$$

$$C_1 = \{\lambda \in C \mid \alpha \neq 0, E \in (-|\alpha|, |\alpha|)\}, \quad (9)$$

$$C_2 = \{\lambda \in C \mid \alpha \neq 0, E \in (|\alpha|, +\infty)\}, \quad (10)$$

$$C_3 = \{\lambda \in C \mid \alpha \neq 0, E = |\alpha|, c \neq 0\}, \quad (11)$$

$$C_4 = \{\lambda \in C \mid \alpha \neq 0, E = -|\alpha|\}, \quad (12)$$

$$C_5 = \{\lambda \in C \mid \alpha \neq 0, E = |\alpha|, c = 0\}, \quad (13)$$

$$C_6 = \{\lambda \in C \mid \alpha = 0, c \neq 0\}, \quad (14)$$

$$C_7 = \{\lambda \in C \mid \alpha = c = 0\}. \quad (14)$$

Further, the sets C_i , $i = 1, \dots, 5$, are divided into subsets determined by the sign of α (see Fig. 1):

$$C_i^+ = C_i \cap \{\alpha > 0\}, \quad C_i^- = C_i \cap \{\alpha < 0\}, \quad i = 1, \dots, 5.$$




FIGURE 1. Stratification of C for $\alpha > 0$ and for $\alpha < 0$

In order to parameterize extremal trajectories, coordinates (φ, k, α) in the domains C_1 and C_2 were introduced in [1] in the following way.

In the domain C_1^+

$$k = \sqrt{\frac{E + \alpha}{2\alpha}} = \sqrt{\frac{c^2}{4\alpha} + \sin^2 \frac{\theta}{2}} \in (0, 1),$$

$$\sin \frac{\theta}{2} = k \operatorname{sn}(\sqrt{\alpha}\varphi), \quad \cos \frac{\theta}{2} = \operatorname{dn}(\sqrt{\alpha}\varphi), \quad \frac{c}{2} = k\sqrt{\alpha} \operatorname{cn}(\sqrt{\alpha}\varphi), \quad \varphi \in [0, 4K(k)].$$

In the domain C_2^+

$$k = \sqrt{\frac{2\alpha}{E + \alpha}} = \frac{1}{\sqrt{\frac{c^2}{4\alpha} + \sin^2 \frac{\theta}{2}}} \in (0, 1),$$

$$\sin \frac{\theta}{2} = \operatorname{sgn} c \operatorname{sn} \frac{\sqrt{\alpha}\varphi}{k}, \quad \cos \frac{\theta}{2} = \operatorname{cn} \frac{\sqrt{\alpha}\varphi}{k}, \quad \frac{c}{2} = \operatorname{sgn} c \frac{\sqrt{\alpha}}{k} \operatorname{dn} \frac{\sqrt{\alpha}\varphi}{k}, \quad \varphi \in [0, 2kK(k)].$$

Here and below dn , sn , cn are Jacobi elliptic functions depending on k , $K(k)$ is the complete elliptic integral of the first kind [27].

In the domains C_1^- , C_2^- the coordinates φ and k are defined as follows:

$$\varphi(\theta, c, \alpha) = \varphi(\theta - \pi, c, -\alpha), \quad k(\theta, c, \alpha) = k(\theta - \pi, c, -\alpha).$$

Immediate differentiation shows that system (7) rectifies in the coordinates (φ, k, α) :

$$\dot{\varphi} = 1, \quad \dot{k} = 0, \quad \dot{\alpha} = 0.$$

In terms of these coordinates, geodesics $q_t = \operatorname{Exp}(\lambda, t)$ with $\lambda = (\theta, c, \alpha) \in \bigcup_{i=1}^3 C_i$ and $\alpha = 1$ are parameterized as follows.

If $\lambda \in C_1$, then

$$\begin{aligned} x_t &= 2k(\operatorname{cn} \varphi_t - \operatorname{cn} \varphi), \\ y_t &= 2(\operatorname{E}(\varphi_t) - \operatorname{E}(\varphi)) - t, \\ z_t &= 2k(\operatorname{sn} \varphi_t \operatorname{dn} \varphi_t - \operatorname{sn} \varphi \operatorname{dn} \varphi - \frac{y_t}{2}(\operatorname{cn} \varphi_t + \operatorname{cn} \varphi)), \\ v_t &= \frac{y_t^3}{6} + 2k^2 \operatorname{cn}^2 \varphi y_t - 4k^2 \operatorname{cn} \varphi (\operatorname{sn} \varphi_t \operatorname{dn} \varphi_t - \operatorname{sn} \varphi \operatorname{dn} \varphi) \\ &\quad + 2k^2 \left(\frac{2}{3} \operatorname{cn} \varphi_t \operatorname{dn} \varphi_t \operatorname{sn} \varphi_t - \frac{2}{3} \operatorname{cn} \varphi \operatorname{dn} \varphi \operatorname{sn} \varphi + \frac{1 - k^2}{3k^2} t + \frac{2k^2 - 1}{3k^2} (\operatorname{E}(\varphi_t) - \operatorname{E}(\varphi)) \right). \end{aligned} \quad (15)$$

Here and below $\operatorname{E}(\varphi) = \int_0^\varphi \operatorname{dn}^2 t dt = E(\operatorname{am}(\varphi), k)$ is the Jacobi epsilon function and $E(u, k)$ is incomplete elliptic integral of the second kind depending on Jacobi amplitude $u = \operatorname{am}(\varphi)$ and k . The Jacobi amplitude $\operatorname{am}(\varphi)$ is the inverse function of the incomplete elliptic integral of the first kind: $F(\operatorname{am}(\varphi)) = \varphi$.

If $\lambda \in C_2$, then

$$\begin{aligned}
x_t &= \frac{2 \operatorname{sgn} c}{k} \left(\operatorname{dn} \psi_t - \operatorname{dn} \psi \right), \\
y_t &= \frac{k^2 - 2}{k^2} t + \frac{2}{k} \left(\operatorname{E}(\psi_t) - \operatorname{E}(\psi) \right), \\
z_t &= -\frac{x_t y_t}{2} - \frac{2 \operatorname{sgn} c \operatorname{dn} \psi}{k} y_t + 2 \operatorname{sgn} c (\operatorname{cn} \psi_t \operatorname{sn} \psi_t - \operatorname{cn} \psi \operatorname{sn} \psi), \\
v_t &= \frac{4}{k} \left(\frac{1}{3} \operatorname{cn} \psi_t \operatorname{dn} \psi_t \operatorname{sn} \psi_t - \frac{1}{3} \operatorname{cn} \psi \operatorname{dn} \psi \operatorname{sn} \psi - \frac{1 - k^2}{3k^3} t - \frac{k^2 - 2}{6k^2} \left(\operatorname{E}(\psi_t) - \operatorname{E}(\psi) \right) \right) \\
&\quad + \frac{y_t^3}{6} + \frac{2y_t}{k^2} \operatorname{dn}^2 \psi - \frac{4}{k} \operatorname{dn} \psi (\operatorname{cn} \psi_t \operatorname{sn} \psi_t - \operatorname{cn} \psi \operatorname{sn} \psi), \\
\psi &= \frac{\varphi}{k}, \quad \psi_t = \psi + \frac{t}{k}.
\end{aligned} \tag{16}$$

If $\lambda \in C_3$, then

$$\begin{aligned}
x_t &= 2 \operatorname{sgn} c \left(\frac{1}{\cosh \varphi_t} - \frac{1}{\cosh \varphi} \right), \\
y_t &= 2(\tanh \varphi_t - \tanh \varphi) - t, \\
z_t &= -\frac{x_t y_t}{2} - \frac{2 \operatorname{sgn} c}{\cosh \varphi} y_t + 2 \operatorname{sgn} c \left(\frac{\tanh \varphi_t}{\cosh \varphi_t} - \frac{\tanh \varphi}{\cosh \varphi} \right), \\
v_t &= \frac{2}{3} \left(\tanh \varphi_t - \tanh \varphi + 2 \frac{\tanh \varphi_t}{\cosh^2 \varphi_t} - 2 \frac{\tanh \varphi}{\cosh^2 \varphi} \right) + \frac{y_t^3}{6} + \frac{2y_t}{\cosh^2 \varphi} - \frac{4}{\cosh \varphi} \left(\frac{\tanh \varphi_t}{\cosh \varphi_t} - \frac{\tanh \varphi}{\cosh \varphi} \right).
\end{aligned} \tag{17}$$

Parameterization of geodesics for $\lambda \in \bigcup_{i=1}^3 C_i$ and arbitrary $\alpha \neq 0$ is obtained from the above parameterization for $\alpha = 1$ via the following symmetries of the Hamiltonian system: dilations

$$\begin{aligned}
\delta_\mu &: (\theta, c, \alpha, t, x, y, z, v) \mapsto (\theta, c/\mu, \alpha/\mu^2, \mu t, \mu x, \mu y, \mu^2 z, \mu^3 v), \quad \mu > 0, \\
\delta_\mu &: (\varphi, k, \alpha) \mapsto (\mu \varphi, k, \alpha/\mu^2),
\end{aligned}$$

and reflection

$$\begin{aligned}
(\theta, c, \alpha, t, x, y, z, v) &\mapsto (\theta - \pi, c, -\alpha, t, -x, -y, z, -v), \\
(\varphi, k, \alpha) &\mapsto (\varphi, k, -\alpha).
\end{aligned}$$

In the remaining cases $\lambda \in \bigcup_{i=4}^7 C_i$ geodesics are parameterized by elementary functions as follows.
 $\lambda \in C_4$:

$$x_t = 0, \quad y_t = t \operatorname{sgn} \alpha, \quad z_t = 0, \quad v_t = \frac{t^3}{6} \operatorname{sgn} \alpha. \tag{18}$$

$\lambda \in C_5$:

$$x_t = 0, \quad y_t = -t \operatorname{sgn} \alpha, \quad z_t = 0, \quad v_t = -\frac{t^3}{6} \operatorname{sgn} \alpha. \tag{19}$$

$\lambda \in C_6 :$

$$\begin{aligned} x_t &= \frac{\cos(ct + \theta) - \cos \theta}{c}, & y_t &= \frac{\sin(ct + \theta) - \sin \theta}{c}, \\ z_t &= \frac{ct - \sin(ct)}{2c^2}, & v_t &= \frac{3 \cos \theta - 2ct \sin \theta - 4 \cos(ct + \theta) + \cos(2ct + \theta)}{4c^3}. \end{aligned} \quad (20)$$

$\lambda \in C_7 :$

$$x_t = -t \sin \theta, \quad y_t = t \cos \theta, \quad z_t = 0, \quad v_t = \frac{t^3}{6} \cos \theta. \quad (21)$$

Projections of geodesics to the plane (x, y) are Euler elasticae (stationary configurations of planar elastic rod with fixed endpoints and tangents at endpoints) [14–16, 28, 29]: inflexional ones for $\lambda \in C_1$, non-inflexional ones for $\lambda \in C_2$, critical ones for $\lambda \in C_3$, straight lines for $\lambda \in C_4 \cup C_5 \cup C_7$, and circles for $\lambda \in C_6$.

1.3. Symmetries of exponential mapping

A pair of mappings

$$s: N \rightarrow N, \quad s: M \rightarrow M$$

is called a symmetry of the exponential mapping if it commutes with this mapping:

$$s \circ \text{Exp}(\lambda, t) = \text{Exp} \circ s(\lambda, t), \quad (\lambda, t) \in N.$$

1.4. Dilations

A one-parameter group of symmetries of the exponential mapping is formed by dilations

$$\begin{aligned} \delta_\mu : (\theta, c, \alpha, t) &\mapsto (\theta, c/\mu, \alpha/\mu^2, \mu t), \\ \delta_\mu : (x, y, z, v) &\mapsto (\mu x, \mu y, \mu^2 z, \mu^3 v), \quad \mu > 0. \end{aligned} \quad (22)$$

1.5. Reflections

The following mappings $\varepsilon^i: C \rightarrow C$ preserve the field of directions of the vertical part of the Hamiltonian vector field $\vec{H}_v = c \frac{\partial}{\partial \theta} - \alpha \sin \theta \frac{\partial}{\partial c} \in \text{Vec}(C)$:

$$\begin{aligned} \varepsilon^1 : (\theta, c, \alpha) &\mapsto (\theta, -c, \alpha), & \varepsilon^2 : (\theta, c, \alpha) &\mapsto (-\theta, c, \alpha), \\ \varepsilon^3 : (\theta, c, \alpha) &\mapsto (-\theta, -c, \alpha), & \varepsilon^4 : (\theta, c, \alpha) &\mapsto (\theta + \pi, c, -\alpha), \\ \varepsilon^5 : (\theta, c, \alpha) &\mapsto (\theta + \pi, -c, -\alpha), & \varepsilon^6 : (\theta, c, \alpha) &\mapsto (-\theta + \pi, c, -\alpha), \\ \varepsilon^7 : (\theta, c, \alpha) &\mapsto (-\theta + \pi, -c, -\alpha). \end{aligned}$$

More precisely, $\varepsilon_*^i \vec{H}_v = \vec{H}_v$ for $i = 3, 4, 7$, and $\varepsilon_*^i \vec{H}_v = -\vec{H}_v$ for $i = 1, 2, 5, 6$. The action of reflections ε^i is continued to symmetries of the exponential mapping as follows.

The action $\varepsilon^i: N \rightarrow N$ is defined as

$$\varepsilon^i(\lambda, t) = \begin{cases} (\varepsilon^i(\lambda), t), & \text{if } \varepsilon_*^i \vec{H}_v = \vec{H}_v, \\ (\varepsilon^i \circ e^{t\vec{H}_v}(\lambda), t), & \text{if } \varepsilon_*^i \vec{H}_v = -\vec{H}_v. \end{cases}$$

The action $\varepsilon^i: M \rightarrow M$ is defined as

$$\varepsilon^i(q) = \varepsilon^i(x, y, z, v) = q^i = (x^i, y^i, z^i, v^i), \quad (23)$$

$$(x^1, y^1, z^1, v^1) = (x, y, -z, v - xz), \quad (24)$$

$$(x^2, y^2, z^2, v^2) = (-x, y, z, v - xz), \quad (25)$$

$$(x^3, y^3, z^3, v^3) = (-x, y, -z, v), \quad (26)$$

$$(x^4, y^4, z^4, v^4) = (-x, -y, z, -v), \quad (27)$$

$$(x^5, y^5, z^5, v^5) = (-x, -y, -z, -v + xz), \quad (28)$$

$$(x^6, y^6, z^6, v^6) = (x, -y, z, -v + xz), \quad (29)$$

$$(x^7, y^7, z^7, v^7) = (x, -y, -z, -v). \quad (30)$$

Thus defined reflections $\varepsilon^i, i = 1, \dots, 7$, form a discrete group of symmetries of the exponential mapping (together with the identity mapping).

1.6. Maxwell points

A point q_t of an extremal trajectory $q_s = \text{Exp}(\lambda, s)$ is called a Maxwell point if there exists another extremal trajectory $\tilde{q}_s = \text{Exp}(\tilde{\lambda}, s)$, $\tilde{q}_s \neq q_s$, such that $\tilde{q}_t = q_t$. The instant t is called a Maxwell time. It is known [10] that an extremal trajectory cannot be optimal after a Maxwell time.

The main result of paper [1], given by Th. 1.1 below, provides an upper bound of the cut time along extremal curves

$$t_{\text{cut}}(\lambda) = \sup\{t > 0 \mid \text{Exp}(\lambda, s) \text{ is optimal for } s \in [0, t]\}.$$

Define the following function $t_{\text{MAX}}^1: C \rightarrow (0, +\infty]$:

$$\lambda \in C_1 \quad \Rightarrow \quad t_{\text{MAX}}^1 = \min(2p_z^1(k), 4K(k))/\sigma, \quad (31)$$

$$\lambda \in C_2 \quad \Rightarrow \quad t_{\text{MAX}}^1 = 2kK(k)/\sigma, \quad (32)$$

$$\lambda \in C_6 \quad \Rightarrow \quad t_{\text{MAX}}^1 = 2\pi/|c|, \quad (33)$$

$$\lambda \in C_3 \cup C_4 \cup C_5 \cup C_7 \quad \Rightarrow \quad t_{\text{MAX}}^1 = +\infty. \quad (34)$$

where $\sigma = \sqrt{|\alpha|}$; $K(k) = \int_0^{\frac{\pi}{2}} \frac{dt}{\sqrt{1 - k^2 \sin^2 t}}$; $p_z^1(k) \in (K(k), 3K(k))$ is the first positive root of the function $f_z(p, k) = \text{dn } p \text{ sn } p + (p - 2E(p)) \text{ cn } p$;

Theorem 1.1 ([1], Th. 3). *For any $\lambda \in C$*

$$t_{\text{cut}}(\lambda) \leq t_{\text{MAX}}^1(\lambda). \quad (35)$$

Proposition 1.2. *The function $t_{\text{MAX}}^1: C \rightarrow (0, +\infty]$ has the following invariant properties:*

- (1) $t_{\text{MAX}}^1(\lambda)$ depends only on the values of E and $|\alpha|$,
- (2) $t_{\text{MAX}}^1(\lambda)$ is an integral of the vector field \vec{H}_v ,
- (3) $t_{\text{MAX}}^1(\lambda)$ is invariant w.r.t. reflections: if $\lambda \in C$, $\lambda^i = \varepsilon^i(\lambda) \in C$, then $t_{\text{MAX}}^1(\lambda^i) = t_{\text{MAX}}^1(\lambda)$,
- (4) t_{MAX}^1 respects the action of dilations: if $\lambda \in C$, $\lambda_\mu = \delta_\mu(\lambda)$, then $t_{\text{MAX}}^1(\lambda_\mu) = \mu t_{\text{MAX}}^1(\lambda)$.

Proof. (1) We denote by \sqcup the union of disjoint sets. Notice first that the decomposition

$$C = C_1 \sqcup C_2 \sqcup C_{35} \sqcup C_4 \sqcup C_6 \sqcup C_7 \quad (36)$$

with $C_{35} = C_3 \cup C_5 = \{\lambda \in C \mid \alpha \neq 0, E = |\alpha|\}$ is determined only by the functions E and $|\alpha|$, see definitions (8)–(14). Thus it remains to show that restriction of t_{MAX}^1 to each of the subsets in decomposition (36) depends only on E and $|\alpha|$.

If $\lambda \in C_1$, then $k = \sqrt{\frac{E + |\alpha|}{2|\alpha|}}$, thus $k = k(E, |\alpha|)$, so $t_{\text{MAX}}^1 = t_{\text{MAX}}^1(E, |\alpha|)$.

The case $\lambda \in C_2$ is similar to the case $\lambda \in C_1$.

If $\lambda \in C_{35} \cup C_4 \cup C_7$, then $t_{\text{MAX}}^1 = +\infty = t_{\text{MAX}}^1(E, |\alpha|)$.

Finally, if $\lambda \in C_6$, then $t_{\text{MAX}}^1 = \frac{2\pi}{|c|} = \frac{\sqrt{2}\pi}{\sqrt{E}}$.

(2) Since E and α are integrals of the vector field \vec{H}_v , then $t_{\text{MAX}}^1 = t_{\text{MAX}}^1(E, |\alpha|)$ is an integral of \vec{H}_v as well.

(3) Let $\lambda \in C$, $\varepsilon^i(\lambda) = \lambda^i \in C$. Since $E(\lambda^i) = E(\lambda)$, $\alpha(\lambda^i) = \pm\alpha(\lambda)$ and $t_{\text{MAX}}^1 = t_{\text{MAX}}^1(E, |\alpha|)$, then $t_{\text{MAX}}^1(\lambda^i) = t_{\text{MAX}}^1(\lambda)$.

(4) Let $\lambda \in C$ and $\lambda_\mu = \delta_\mu(\lambda)$, $\mu > 0$. Since we have $E(\lambda_\mu) = \frac{1}{\mu^2}E(\lambda)$ and $\alpha(\lambda_\mu) = \frac{1}{\mu^2}\alpha(\lambda)$, then $\delta_\mu(C_i) = C_i$, $i = 1, \dots, 7$, and $k(\lambda_\mu) = k(\lambda)$. Then it follows from the definition of the function t_{MAX}^1 that $t_{\text{MAX}}^1(\lambda_\mu) = \mu \cdot t_{\text{MAX}}^1(\lambda)$ for $\lambda \in C_i$ and each $i = 1, \dots, 7$. \square

1.7. Conjugate points

A point $q_t = \text{Exp}(\lambda, t)$ is called a conjugate point for q_0 if $\nu = (\lambda, t)$ is a critical point of the exponential mapping and that is why q_t is the corresponding critical value:

$$d_\nu \text{Exp}: T_\nu N \rightarrow T_{q_t} M \text{ is degenerate.}$$

The instant t is called a conjugate time along the extremal trajectory $q_s = \text{Exp}(\lambda, s)$, $s \geq 0$.

The first conjugate time along a trajectory $\text{Exp}(\lambda, s)$ is denoted by

$$t_{\text{conj}}^1(\lambda) = \min \{t > 0 \mid t \text{ is a conjugate time along } \text{Exp}(\lambda, s), s \geq 0\}.$$

The trajectory $\text{Exp}(\lambda, s)$ loses its local optimality at the instant $t = t_{\text{conj}}^1(\lambda)$ (see [4]).

The following lower bound on the first conjugate time is the main result of work [2].

Theorem 1.3 ([2]). *For any $\lambda \in C$*

$$t_{\text{conj}}^1(\lambda) \geq t_{\text{MAX}}^1(\lambda). \quad (37)$$

2. DECOMPOSITIONS IN PREIMAGE AND IMAGE OF EXPONENTIAL MAPPING

In this section we describe decomposition (43) in the image, and decomposition (49) in the preimage of the exponential mapping, which will be proved to be diffeomorphic via the exponential mapping in Th. 3.20.

2.1. Decomposition in M

Let $\widehat{M} = M \setminus \{q_0\}$, then $M = \widehat{M} \sqcup \{q_0\}$. Further, we denote the subset containing the Maxwell strata MAX^1 and MAX^2 :

$$M' = \{q \in \widehat{M} \mid xz = 0\}$$

and its complement

$$\widetilde{M} = \{q \in \widehat{M} \mid xz \neq 0\},$$

then

$$\widehat{M} = \widetilde{M} \sqcup M'. \quad (38)$$

Denote the connected components of the set \widetilde{M} :

$$M_1 = \{q \in M \mid x < 0, z > 0\}, \quad (39)$$

$$M_2 = \{q \in M \mid x < 0, z < 0\}, \quad (40)$$

$$M_3 = \{q \in M \mid x > 0, z < 0\}, \quad (41)$$

$$M_4 = \{q \in M \mid x > 0, z > 0\}, \quad (42)$$

so that

$$\widetilde{M} = \bigsqcup_{i=1}^4 M_i. \quad (43)$$

This decomposition agrees with the action of reflections and dilations as described in the following statement.

Proposition 2.1.

- (1) Reflections $\varepsilon^j \in G$ permute the domains M_i according to Table 1.
- (2) Dilations δ_μ , $\mu > 0$, preserve the domains M_i .

Id, ε^6	$\varepsilon^1, \varepsilon^7$	$\varepsilon^2, \varepsilon^4$	$\varepsilon^3, \varepsilon^5$
M_1	M_2	M_4	M_3
M_2	M_1	M_3	M_4
M_3	M_4	M_2	M_1
M_4	M_3	M_1	M_2

TABLE 1. Action of the reflections ε^j on the domains M_i

Proof. Follows immediately from the definitions of the actions of reflections $\varepsilon^j: M \rightarrow M$, see (23)–(30), and dilations $\delta_\mu: M \rightarrow M$, see (22). \square

2.2. Decomposition in N

Denote the subset in preimage of the exponential mapping that corresponds to all potentially optimal geodesics:

$$\widehat{N} = \{(\lambda, t) \in N \mid t \leq t_{\text{MAX}}^1(\lambda)\}.$$

If $(\lambda, t) \in N \setminus \widehat{N}$, then the geodesic $\text{Exp}(\lambda, s)$, $s \in [0, t]$, is non-optimal. We decompose the set \widehat{N} into subsets corresponding to the subsets of the set \widehat{M} (Subsec. 2.1), the proof of this correspondence will be given in Subsec. 2.3. Let

$$N' = \{(\lambda, t) \in N \mid t = t_{\text{MAX}}^1(\lambda) \text{ or } c_{t/2} \sin \theta_{t/2} = 0\},$$

$$\widetilde{N} = \{(\lambda, t) \in N \mid t < t_{\text{MAX}}^1(\lambda), c_{t/2} \sin \theta_{t/2} \neq 0\},$$

then

$$\widehat{N} = \widetilde{N} \sqcup N'. \quad (44)$$




FIGURE 2. The trace of domains D_i in the set $\{t = 0\}$ for $\alpha \geq 0$ and for $\alpha < 0$

The following sets will play an important role in the description of the global structure of the exponential mapping:

$$D_1 = \{(\lambda, t) \in N \mid t \in (0, t_{\text{MAX}}^1(\lambda)), \sin \theta_{t/2} > 0, c_{t/2} > 0\}, \quad (45)$$

$$D_2 = \{(\lambda, t) \in N \mid t \in (0, t_{\text{MAX}}^1(\lambda)), \sin \theta_{t/2} > 0, c_{t/2} < 0\}, \quad (46)$$

$$D_3 = \{(\lambda, t) \in N \mid t \in (0, t_{\text{MAX}}^1(\lambda)), \sin \theta_{t/2} < 0, c_{t/2} < 0\}, \quad (47)$$

$$D_4 = \{(\lambda, t) \in N \mid t \in (0, t_{\text{MAX}}^1(\lambda)), \sin \theta_{t/2} < 0, c_{t/2} > 0\}. \quad (48)$$

We have the obvious decomposition

$$\tilde{N} = \bigsqcup_{i=1}^4 D_i. \quad (49)$$

The trace of domains D_i in the set $\{(\lambda, t) \in N \mid t = 0\}$ is shown in Fig. 2.

Proposition 2.2.

- (1) Reflections $\varepsilon^j \in G$ permute the sets D_i according to Table 2.
- (2) Dilations $\delta_\mu, \mu > 0$, preserve the sets D_i .

Id, ε^6	$\varepsilon^1, \varepsilon^7$	$\varepsilon^2, \varepsilon^4$	$\varepsilon^3, \varepsilon^5$
D_1	D_2	D_4	D_3
D_2	D_1	D_3	D_4
D_3	D_4	D_2	D_1
D_4	D_3	D_1	D_2

TABLE 2. Action of the reflections ε^j on the domains D_i

Proof. (1) We prove only the equality $\varepsilon^1(D_1) = D_2$, all the rest equalities given in Table 2 are proved similarly. Let $(\lambda, t) = (\theta, c, \alpha, t) \in D_1$ and $\varepsilon^1(\lambda, t) = (\lambda^1, t) = (\theta^1, c^1, \alpha^1, t)$, we show that $(\lambda^1, t) \in D_2$.

Denote $\lambda_{t/2} = (\theta_{t/2}, c_{t/2}, \alpha) = e^{(t/2)\vec{H}_v}(\lambda)$ and $\lambda_{t/2}^1 = (\theta_{t/2}^1, c_{t/2}^1, \alpha^1) = e^{(t/2)\vec{H}_v}(\lambda^1)$. Since $\varepsilon_*^1 \vec{H}_v = -\vec{H}_v$, then $\lambda^1 = \varepsilon^1 \circ e^{t\vec{H}_v}(\lambda)$, thus

$$\lambda_{t/2}^1 = e^{(t/2)\vec{H}_v} \circ \varepsilon^1 \circ e^{t\vec{H}_v}(\lambda) = \varepsilon^1 \circ e^{-(t/2)\vec{H}_v} \circ e^{t\vec{H}_v}(\lambda) = \varepsilon^1 \circ e^{(t/2)\vec{H}_v}(\lambda) = \varepsilon^1(\lambda_{t/2}).$$

That is,

$$(\theta_{t/2}^1, c_{t/2}^1, \alpha^1) = (\theta_{t/2}, -c_{t/2}, \alpha).$$

The inclusion $(\lambda, t) \in D_1$ means that

$$t \in (0, t_{\text{MAX}}^1(\lambda)), \quad \sin \theta_{t/2} > 0, \quad c_{t/2} > 0,$$

thus $\sin \theta_{t/2}^1 > 0$, $c_{t/2}^1 < 0$. Moreover, since $t_{\text{MAX}}^1(\lambda^1) = t_{\text{MAX}}^1 \circ \varepsilon^1(\lambda) = t_{\text{MAX}}^1(\lambda)$ by Propos. 1.2, then $t \in (0, t_{\text{MAX}}^1(\lambda^1))$. Consequently, $(\lambda^1, t) \in D_2$.

We proved that $\varepsilon^1(D_1) \subset D_2$. It follows similarly that $\varepsilon^1(D_2) \subset D_1$. Since $\varepsilon^1 \circ \varepsilon^1 = \text{Id}$ on N , we have $\varepsilon^1(D_1) = D_2$.

(2) Let $(\lambda, t) = (\theta, c, \alpha, t) \in N$, $(\lambda_\mu, t_\mu) = \delta_\mu(\lambda, t) = (\theta, c/\mu, \alpha/\mu^2, \mu t)$. Since $t_{\text{MAX}}^1(\lambda_\mu) = \mu t_{\text{MAX}}^1(\lambda)$ by Propos. 1.2, it is obvious that if $(\lambda, t) \in D_1$, then $(\lambda_\mu, t_\mu) \in D_1$. Thus $\delta_\mu(D_1) \subset D_1$. Since $d_{1/\mu} = (\delta_\mu)^{-1}$, then $\delta_\mu(D_1) = D_1$. It follows similarly that $\delta_\mu(D_i) = D_i$ for $i = 2, 3, 4$. \square

2.3. Basic properties of exponential mapping

In this subsection we describe some simple properties on the action of the exponential mapping on the subsets of N defined in the previous subsection.

First of all, $\text{Exp}(\widehat{N}) \supset \widehat{M}$ since for any point $q_1 \in \widehat{M}$ there exists an optimal trajectory $q_s = \text{Exp}(\lambda, s)$ such that $q_{t_1} = q_1$, thus $t_1 \leq t_{\text{cut}}(\lambda) \leq t_{\text{MAX}}^1(\lambda)$, i.e., $\text{Exp}(\lambda, t_1) = q_1$ with $(\lambda, t_1) \in \widehat{N}$. However, Maxwell points in \widehat{M} have several preimages in \widehat{N} . Moreover, the mapping $\text{Exp}|_{\widehat{N}}$ is degenerate at points (λ, t) where $t = t_{\text{MAX}}^1(\lambda)$ is a conjugate time along the trajectory $\text{Exp}(\lambda, s)$.

In the next two propositions we show that the action of Exp is compatible with decompositions (44), (38), and (49), (43).

Proposition 2.3. *There holds the inclusion*

$$\text{Exp}(N') \subset \{q \in M \mid xz = 0\} = M' \sqcup \{q_0\}. \quad (50)$$

Proof. The reflection ε^4 and dilations δ_μ , $\mu > 0$, are symmetries of Exp and preserve the sets N' , M' and $\{q_0\}$. Since $\varepsilon^4: \alpha \mapsto -\alpha$ and $\delta_\mu: \alpha \mapsto \alpha/\mu^2$, we can assume in the proof of inclusion (50) that $\alpha \in \{0, 1\}$.

Let $(\lambda, t) \in N'$ and $q_t = (x_t, y_t, z_t, v_t) = \text{Exp}(\lambda, t)$, we show that $x_t z_t = 0$.

Suppose first that $\alpha = 1$, then $\lambda \in \bigcup_{i=1}^5 C_i$.

Let $\lambda \in C_1$, then we use parameterization of extremals (15). Since $(\lambda, t) \in N'$, then $c_{t/2} \sin \theta_{t/2} = 0$ or $t = t_{\text{MAX}}^1(\lambda)$. If $c_{t/2} = 2k \text{cn} \tau = 0$, then $\text{cn} \tau = 0$, thus $z_t = 0$ in view of (7.3) [1]. If $\sin \theta_{t/2} = 2k \text{sn} \tau \text{dn} \tau = 0$, then $x_t = 0$ in view of (7.2) [1]. Finally, if $t = t_{\text{MAX}}^1(\lambda)$, then $p = p_z^1(k)$ or $p = 2K(k)$ by (31), thus $z_t = 0$ or $x_t = 0$ by (7.2) and (7.3) [1].

The case $\lambda \in C_2 \cup C_3$ is considered similarly to the case $\lambda \in C_1$.

If $\lambda \in C_4 \cup C_5$, then $x_t = 0$ by (18), (19).

Now suppose that $\alpha = 0$, thus $\lambda \in C_6 \cup C_7$.

Let $\lambda \in C_6$, then we use parameterization of extremals (20). The case $c_{t/2} = 0$ is impossible. If $\sin \theta_{t/2} = 0$, then $t = 2\pi/|c|$, thus $x_t = 0$. If $t = t_{\text{MAX}}^1(\lambda) = 2\pi/|c|$ (33), then $x_t = 0$ as well.

Finally, if $\lambda \in C_7$, then $z_t = 0$ by (20). \square

Proposition 2.4. *For any $i = 1, \dots, 4$, we have $\text{Exp}(D_i) \subset M_i$.*

Proof. By virtue of the reflections ε^i (Propos. 2.1, 2.2), the proof of this proposition reduces to the case $i = 1$. So let $(\lambda, t) \in D_1$, we prove that $q_t = \text{Exp}(\lambda, t) \in M_1$.

The reflection ε^6 and the dilations $\delta_\mu, \mu > 0$, preserve the domains D_1 and M_1 , and act on the parameter α as $\varepsilon^6 : \alpha \mapsto -\alpha$, $\delta_\mu : \alpha \mapsto \frac{\alpha}{\mu^2}$, thus we can assume in this proof that $\alpha \in \{0, 1\}$.

Since $(\lambda, t) \in D_1$, then $\sin \theta_{t/2} > 0$, $c_{t/2} > 0$, $t \in (0, t_{\text{MAX}}^1(\lambda))$.

Let $\alpha = 1$, then $\lambda \in C_1 \cup C_2 \cup C_3$.

Let $\lambda \in C_1$. Then $\sin \theta_{t/2} = 2k \text{sn} \tau \text{dn} \tau > 0$, $c_{t/2} = 2k \text{cn} \tau > 0$. Since in this case $t \in (0, t_{\text{MAX}}^1(\lambda))$ and $t_{\text{MAX}}^1(\lambda) = \min(2p_z^1, 4K)$, then $f_z(p, k) > 0$ and $\text{sn} p > 0$. Then formulas (7.2), (7.3) [1] imply that $x_t < 0, z_t > 0$, i.e., $q_t \in M_1$.

The cases $\lambda \in C_2$ and $\lambda \in C_3$ are considered similarly to the case $\lambda \in C_1$.

Now let $\alpha = 0$, then $\lambda \in C_6$. Then $x_t = -\frac{2}{c} \sin \theta_{t/2} \sin \frac{ct}{2} < 0$ and $z_t = \frac{ct - \sin(ct)}{2c^2} > 0$, thus $q_t \in M_1$.

We proved that $\text{Exp}(D_1) \subset M_1$. □

Our goal is to prove that the mappings $\text{Exp}: D_i \rightarrow M_i$, $i = 1, \dots, 4$, are diffeomorphisms, see Th. 3.21. This is done in Section 3 via the following Hadamard global diffeomorphism theorem.

Theorem 2.5 ([18]). *Let $F: X \rightarrow Y$ be a smooth mapping between smooth manifolds of equal dimension. Let the following conditions hold:*

- (1) X is connected,
- (2) Y is connected and simply connected,
- (3) F is nondegenerate,
- (4) F is proper (i.e., $F^{-1}(K) \subset X$ is compact for a compact $K \subset Y$).

Then F is a diffeomorphism.

2.4. Topological properties of decompositions in M and N

We prove that hypotheses (1), (2) of Th. 2.5 are verified for the mappings $\text{Exp}: D_i \rightarrow M_i$.

Definition 2.6. Suppose X is a topological space and $f_1, f_2: X \rightarrow \mathbb{R}$. Then $f_1 \sim f_2$ on a sequence $\{\lambda_n\} \subset X$ if $\lim_{n \rightarrow \infty} \frac{f_1(\lambda_n)}{f_2(\lambda_n)} = 1$.

Proposition 2.7.

- (1) The sets $D_i \subset N$, $i = 1, \dots, 4$, are open and connected.
- (2) The sets $M_i \subset M$, $i = 1, \dots, 4$, are open, connected and simply connected.

In the proof of item (1) of this proposition we need the following statement.

Proposition 2.8. *The function $t_{\text{MAX}}^1: C \rightarrow (0, +\infty]$ is continuous on the set $C \setminus C_4$, and is smooth on the set $C_1^0 \cup C_2$, where $C_1^0 = \{\lambda \in C_1 \mid k \neq k_0\}$.*

Remark 2.9. We assume in $(0, +\infty]$ the natural basis of topology:

$$(a, b), \quad (a, +\infty], \quad 0 < a < b < +\infty.$$

Proof. Let $\lambda_n \rightarrow \bar{\lambda}$ as $n \rightarrow \infty$, where $\lambda_n, \bar{\lambda} \in C \setminus C_4 = (\cup_{i=1}^3 C_i) \cup (\cup_{i=5}^7 C_i)$. We denote $t_n = t_{\text{MAX}}^1(\lambda_n)$ and $\bar{t} = t_{\text{MAX}}^1(\bar{\lambda})$, then prove that $t_n \rightarrow \bar{t}$ as $n \rightarrow +\infty$.

1. Let $\lambda_n \in C_1$, then $\bar{\lambda} \in \text{cl}(C_1) \setminus C_4 = C_1 \cup C_3 \cup C_5 \cup C_7$.

1.1. Let $\bar{\lambda} \in C_1$. The function $t_{\text{MAX}}^1|_{C_1} = \frac{\min(2p_z^1(k), 4K(k))}{\sigma}$ is continuous since for $k \in (0, 1)$ the function $\min(p_z^1(k), 2K(k))$ is continuous (see Cor. 3.1 [16]), thus $t_n \rightarrow \bar{t}$.

1.2. Let $\bar{\lambda} \in C_3 \cup C_5$. Then $k_n = k(\lambda_n) \rightarrow 1$, $K(k_n) \rightarrow +\infty$, $p_z^1(k_n) \rightarrow +\infty$, $\sigma(\lambda_n) \rightarrow \bar{\sigma} > 0$. Thus $t_n \rightarrow +\infty = \bar{t}$.

1.3. Let $\bar{\lambda} \in C_7$. Then $\alpha_n \rightarrow 0$. Since $\min(2p_z^1(k), 4K(k)) > 2K(k) > \pi$, then $t_n \rightarrow +\infty = \bar{t}$.

2. Let $\lambda_n \in C_2$, then $\bar{\lambda} \in \text{cl}(C_2) \setminus C_4 = C_2 \cup C_3 \cup C_5 \cup C_6 \cup C_7$.

2.1. Let $\bar{\lambda} \in C_2$. The function $t_{\text{MAX}}^1|_{C_2} = \frac{2K(k)k}{\sigma}$ is continuous, thus $t_n \rightarrow \bar{t}$.

2.2. Let $\bar{\lambda} \in C_3 \cup C_5$. This case is similar to case 1.2.

2.3. Let $\bar{\lambda} \in C_6$. Then

$$\begin{aligned} \alpha_n &= \alpha(\lambda_n) \rightarrow \bar{\alpha} = 0, & c_n &= c(\lambda_n) \rightarrow \bar{c} \neq 0, \\ E_n &= E(\lambda_n) \rightarrow \frac{\bar{c}^2}{2} = \bar{E} \neq 0, & k_n &= k(\lambda_n) = \sqrt{\frac{2|\alpha_n|}{E_n + |\alpha_n|}} \sim \frac{2\sqrt{|\alpha_n|}}{|\bar{c}|} \rightarrow 0, \\ t_n &= \frac{2K(k_n)k_n}{\sqrt{|\alpha_n|}} \sim 2K(0) \cdot \frac{2}{|\bar{c}|} = \frac{2\pi}{|\bar{c}|} = \bar{t}, \end{aligned}$$

i.e., $t_n \rightarrow \bar{t}$.

2.4. Let $\bar{\lambda} \in C_7$. Then $\alpha_n \rightarrow 0, c_n \rightarrow 0$. Thus $E_n \rightarrow 0$, so $\frac{k_n}{\sqrt{|\alpha_n|}} = \sqrt{\frac{2}{E_n + |\alpha_n|}} \rightarrow +\infty$. Consequently, $t_n = \frac{2K(k_n)k_n}{\sqrt{|\alpha_n|}} \rightarrow +\infty = \bar{t}$.

3. Let $\lambda_n \in C_3$, then $\bar{\lambda} \in \text{cl}(C_3) \setminus C_4 = C_3 \cup C_5 \cup C_7$, and this case is similar to cases 1.2 and 1.3.

4. Let $\lambda_n \in C_5$, then $\bar{\lambda} \in \text{cl}(C_5) = C_5 \cup C_7$, and $t_n = +\infty = \bar{t}$.

5. Let $\lambda_n \in C_6$, then $\bar{\lambda} \in \text{cl}(C_6) = C_6 \cup C_7$.

5.1. Let $\bar{\lambda} \in C_6$. Since the function $t_{\text{MAX}}^1|_{C_6} = \frac{2\pi}{|c|}$ is continuous, then $t_n \rightarrow \bar{t}$.

5.2. Let $\bar{\lambda} \in C_7$. Then $c_n \rightarrow 0$, thus $t_n = \frac{2\pi}{|c_n|} \rightarrow +\infty = \bar{t}$.

The function $t_{\text{MAX}}^1(\lambda)$ is smooth on C_1^0 since for $\lambda \in C_1$ we have by virtue of (31):

$$\begin{aligned} k < k_0 &\Rightarrow t_{\text{MAX}}^1 = \frac{2p_z^1(k)}{\sqrt{|\alpha|}} \in C^\infty, \\ k > k_0 &\Rightarrow t_{\text{MAX}}^1 = \frac{4K}{\sqrt{|\alpha|}} \in C^\infty. \end{aligned}$$

Similarly, $t_{\text{MAX}}^1(\lambda)$ is smooth on C_2 by virtue of (32). The proof of Propos. 2.8 is complete. \square

Remark 2.10. The function t_{MAX}^1 is discontinuous on C_4 .

Indeed, let $\lambda_n \in C_1$ be such that $k(\lambda_n) \rightarrow 0$ and $\alpha(\lambda_n) \rightarrow \bar{\alpha} \neq 0$. Then $\lambda_n \rightarrow \bar{\lambda} \in C_4$ but

$$t_{\text{MAX}}^1(\lambda_n) \rightarrow \frac{2p_z^1(0)}{\sqrt{|\bar{\alpha}|}} < t_{\text{MAX}}^1(\bar{\lambda}) = +\infty.$$

Here $p = p_z^1(0)$ is the minimal positive root of the equation $f_z(p, 0) = \sin p - p \cos p = 0$, thus $p \in (\pi, 3\pi/2)$.

Now we prove Propos. 2.7.

Proof. (1) Reflections $\varepsilon^i: N \rightarrow N$ are diffeomorphisms and permute the sets D_i , thus it is sufficient to prove that the set $D_1 = \{(\lambda, t) \in N \mid \sin \theta_{t/2} > 0, c_{t/2} > 0, t < t_{\text{MAX}}^1(\lambda)\}$ is open and connected.

Consider the vector field $P = \frac{t}{2}(c \frac{\partial}{\partial \theta} - \alpha \sin \theta \frac{\partial}{\partial c}) \in \text{Vec}(N)$. Denote the flow of this vector field for time 1 as $e^P \in \text{Diff}(N)$. We have

$$e^P(\theta, c, \alpha, t) = e^P(\lambda, t) = (e^{\frac{t}{2}\bar{H}^v}(\lambda), t) = (\theta_{t/2}, c_{t/2}, \alpha, t),$$

thus $e^P(D_1) = \tilde{D}_1$, where

$$\tilde{D}_1 = \{(\lambda, t) \in N \mid \sin \theta > 0, c > 0, t < t_{\text{MAX}}^1(\lambda, t)\}.$$

By Propos. 2.8, the function $(t - t_{\text{MAX}}^1(\lambda)): N \rightarrow (0, +\infty]$ is continuous on the set $N \setminus N_4 \supset \tilde{D}_1$, thus the set \tilde{D}_1 is open. Moreover, the domain \tilde{D}_1 is a subgraph of the function $t_{\text{MAX}}^1(\lambda)$ on a connected domain $\{(\theta, c, \alpha) \in C \mid \theta \in (0, \pi), c > 0, \alpha \in \mathbb{R}\}$, thus \tilde{D}_1 is connected.

We proved that \tilde{D}_1 is open and connected, thus $D_1 = e^{-P}(\tilde{D}_1)$ is open and connected as well.

(2) It is obvious from definitions (39)–(42) that the sets M_i , $i = 1, \dots, 4$, are open, connected and simply connected. \square

3. DIFFEOMORPHIC PROPERTIES OF EXPONENTIAL MAPPING

In this section we prove that restriction of the exponential mapping to the subdomains D_i, M_i is a diffeomorphism.

Lemma 3.1. *If $\text{Exp}: D_1 \rightarrow M_1$ is proper, then $\text{Exp}: D_i \rightarrow M_i$ is proper for $i = 2, 3, 4$.*

Proof. Follows immediately from Propositions 2.1, 2.2. \square

Lemma 3.2. *The mapping $\text{Exp}: D_1 \rightarrow M_1$ is proper iff there exists no sequence $\{\nu_n\} \subset D_1 = (D_1 \cap N_1) \cup (D_1 \cap N_2) \cup (D_1 \cap N_3) \cup (D_1 \cap N_6)$, such that $\nu_n \rightarrow \bar{\nu} \in \text{cl}(D_1) \setminus D_1$ and $\text{Exp}(\nu_n) \rightarrow \bar{q} \in M_1$.*

Proof. It follows from the definition of a proper mapping that the mapping $\text{Exp}: D_1 \rightarrow M_1$ is proper iff there exists no sequence $\{\nu_n\} \subset D_1$, such that $\nu_n \rightarrow \bar{\nu} \in \text{cl}(D_1) \setminus D_1$ and $\text{Exp}(\nu_n) \rightarrow \bar{q} \in M_1$.

Moreover, the definition of D_1 (45) gives the decomposition

$$D_1 = (D_1 \cap N_1) \cup (D_1 \cap N_2) \cup (D_1 \cap N_3) \cup (D_1 \cap N_6).$$

\square

Let us introduce the following sets for arbitrary $\varepsilon \in (0, 1)$:

$$S_\varepsilon := \{\nu \in N \mid \theta_{t/2} \in [\varepsilon, \pi - \varepsilon], c_{t/2} \in [\varepsilon, 1/\varepsilon], |\alpha| \leq 1/\varepsilon, t \in [\varepsilon, 1/\varepsilon], t_{\text{MAX}}^1(\lambda) - t \geq \varepsilon\}.$$

Lemma 3.3. *The set S_ε is compact for any $\varepsilon > 0$.*

Proof. Let $\{\nu_n\} \subset S_\varepsilon$ be an arbitrary sequence. To prove the lemma, we need to find a subsequence ν_{n_m} which tends to $\bar{\nu} \in S_\varepsilon$ as $m \rightarrow \infty$.

Since α, t are bounded on S_ε , we obtain for a subsequence that $\alpha \rightarrow \bar{\alpha}, t \rightarrow \bar{t}$ as $m \rightarrow \infty$.

Since $\theta_{t/2}, c_{t/2}$ are bounded on S_ε , we obtain for a subsequence $(\theta_{t/2}, c_{t/2}) \rightarrow (a, b)$. Moreover, we have $(\theta, c) = \Phi_{-t/2}(\theta_{t/2}, c_{t/2}) \rightarrow \Phi_{-\bar{t}/2}(a, b) =: (\bar{\theta}, \bar{c})$, where Φ is the flow of pendulum (7). Since $\bar{\nu} = (\bar{\theta}, \bar{c}, \bar{\alpha}, \bar{t}) \in S_\varepsilon$ by continuity of the functions which define S_ε , we see that S_ε is compact. \square

Lemma 3.4. *If $K \subset D_1$ is compact, then there exists $\varepsilon > 0$ such that $K \subset S_\varepsilon$.*

Proof. Since the functions $\theta_{t/2}$, $c_{t/2}$, α , t , $(t_{\text{MAX}}^1 - t)$ are continuous on N , these functions attain maximum and minimum on K . \square

Lemma 3.5. *Let $\{\nu_n\} \subset D_1$. Then $\nu_n \rightarrow \bar{\nu} \in \text{cl}(D_1) \setminus D_1$ iff one of the following conditions holds for $\{\nu_n\}$:*

- (1) $\theta_{t/2} \rightarrow 0$,
- (2) $\theta_{t/2} \rightarrow \pi$,
- (3) $c_{t/2} \rightarrow 0$,
- (4) $c_{t/2} \rightarrow +\infty$,
- (5) $t \rightarrow 0$,
- (6) $t_{\text{MAX}}^1(\lambda) - t \rightarrow 0$,
- (7) $|\alpha| \rightarrow \infty$.

Proof. Necessity. Assume the converse. Suppose for any sequence $\{\nu_n\} \subset D_1$, $\nu_n \rightarrow \bar{\nu} \in \text{cl}(D_1) \setminus D_1$, that conditions (1)–(7) do not hold. This means that there exists $\varepsilon > 0$ such that conditions

$$\theta_{t/2} \geq \varepsilon, \theta_{t/2} \leq \pi - \varepsilon, \quad c_{t/2} \geq \varepsilon, c_{t/2} \leq 1/\varepsilon, \quad t \geq \varepsilon, t_{\text{MAX}}^1(\lambda) - t \geq \varepsilon, \quad |\alpha| \leq 1/\varepsilon$$

hold for a subsequence. It follows that $\{\nu_n\} \subset S_\varepsilon$, which is a compact subset of D_1 . So $\bar{\nu} \in S_\varepsilon \subset D_1$. This contradiction proves the necessity.

Sufficiency. Assume the converse. Let for any sequence $\nu_n \subset D_1$ we have $\nu_n \rightarrow \bar{\nu} \in D_1$. Then there exists a compact set $K \supset \{\nu_n\}$, $\bar{\nu} \in K$. This means that there exists $\varepsilon > 0$ such that $K \subset S_\varepsilon$. This contradiction proves the lemma. \square

Definition 3.6. Suppose X is a topological space and $f_1, f_2: X \rightarrow \mathbb{R}$. Then $f_1 \approx f_2$ on a sequence $\{\nu_n\} \subset X$ if $\lim_{n \rightarrow \infty} \frac{f_1(\nu_n)}{f_2(\nu_n)} \in \mathbb{R} \setminus \{0\}$.

In the next lemmas we use the parametrization of exponential mapping for the case $\lambda \in C_6$, see (20).

Lemma 3.7. *If $\{\nu_n\} \subset D_1 \cap N_6$, $\nu_n \rightarrow \bar{\nu} \in \text{cl}(D_1) \setminus D_1$, and $\text{Exp}(\nu_n) \rightarrow \bar{q} \in M_1$, then $c \rightarrow 0$ on the sequence $\{\nu_n\}$.*

Proof. Notice that for $\bar{\nu} = \{\bar{x}, \bar{y}, \bar{z}, \bar{v}\} \in M_1$ we have $\bar{x} \neq 0$. Consider all possible cases $\bar{\nu} \in \text{cl} D_1 \setminus D_1$:

1. $\theta_{t/2} \rightarrow 0 \Rightarrow \frac{ct}{2} + \theta \rightarrow 0 \Rightarrow c \rightarrow 0$ or $x = -\frac{2 \sin(\frac{ct}{2} + \theta) \sin \frac{ct}{2}}{c} \rightarrow \bar{x} = 0$.
2. $\theta_{t/2} \rightarrow \pi \Rightarrow \frac{ct}{2} + \theta \rightarrow 0 \Rightarrow c \rightarrow 0$ or $x \rightarrow \bar{x} = 0$.
3. $c_{t/2} \rightarrow 0 \Rightarrow c \rightarrow 0$.
4. $c_{t/2} \rightarrow \infty \Rightarrow c \rightarrow \infty \Rightarrow x \rightarrow \bar{x} = 0$.
5. $t \rightarrow 0 \Rightarrow c \rightarrow 0$, otherwise $x \rightarrow \bar{x} = 0$.
6. $t \rightarrow \frac{2\pi}{|c|}$. This means that $c \rightarrow 0$ or $x \rightarrow \bar{x} = 0$. \square

Lemma 3.8. *Suppose $\nu_n \in D_1 \cap N_6$. If $c \rightarrow 0$, then $x \rightarrow 0$ or one of the functions x, y, z tends to ∞ on the sequence $\{\nu_n\}$.*

Proof. Consider two possible cases:

1. If $ct \rightarrow 0$, then

$$z \approx \frac{(ct)^3}{c^2} = ct^3,$$

$$x^2 + y^2 = \frac{2 - 2(\cos(ct + \theta) \cos \theta + \sin(ct + \theta) \sin \theta)}{c} = \frac{2(1 - \cos(ct))}{c} \approx ct^2.$$

It follows that $t \rightarrow \infty$, otherwise $x^2 \rightarrow 0$. Then we have $z \approx (x^2 + y^2)t$. This means that $z \approx t \rightarrow \infty$ or $x^2 \rightarrow 0$.

2. If $ct \rightarrow \bar{c} \neq 0$ or $ct \rightarrow \infty$, then $ct - \sin(ct) > M > 0$, thus $z \rightarrow \infty$. \square

In the next lemmas we use the following parametrization of exponential mapping for the case $\lambda \in C_3$ (see (17)):

$$\begin{aligned} x &= -\frac{8 \operatorname{sgn} c \sigma \sinh p \sinh \tau}{\alpha (\cosh(2p) + \cosh(2\tau))}, \\ y &= \frac{2\sigma}{\alpha} \left(\frac{2 \sinh(2p)}{\cosh(2p) + \cosh(2\tau)} - p \right), \\ z &= \frac{8 \operatorname{sgn} c \cosh \tau (p \cosh p - \sinh p)}{|\alpha| (\cosh(2p) + \cosh(2\tau))}, \\ v &= -\frac{1}{3\alpha\sigma \cosh(p-\tau) \cosh^2(p+\tau)} \left(6(\cosh \tau - 3 \cosh(2p+\tau)) \sinh p \right. \\ &\quad \left. + 2p \left(6 \cosh(3p+\tau) + p \cosh(p+\tau) (p(\cosh(2p) + \cosh(2\tau)) - 6 \sinh(2p)) \right) \right). \end{aligned}$$

Lemma 3.9. *If $\{\nu_n\} \subset D_1 \cap N_3$, $\nu_n \rightarrow \bar{\nu} \in \operatorname{cl}(D_1) \setminus D_1$, and $\operatorname{Exp}(\nu_n) \rightarrow \bar{q} \in M_1$, then $\sigma \rightarrow 0$ or $p \rightarrow \infty$ and $t \rightarrow \infty$ with $\sigma \rightarrow \bar{\sigma} \neq 0$.*

Proof. Notice that for $\bar{\nu} = \{\bar{x}, \bar{y}, \bar{z}, \bar{v}\} \in M_1$ we have $\bar{x} \neq 0$. Consider all possible cases $\nu \rightarrow \operatorname{cl}(D_1) \setminus D_1$:

$$1. \theta_{t/2} \rightarrow 0 \Rightarrow \begin{cases} \tanh \tau \rightarrow 0 \\ \cosh \tau \rightarrow 1 \end{cases} \Rightarrow \tau \rightarrow 0 \Rightarrow \sigma \rightarrow 0, \text{ or } x \rightarrow \bar{x} = 0, \text{ or } z \rightarrow \infty \text{ with } p \rightarrow \infty.$$

$$2. \theta_{t/2} \rightarrow \pi \Rightarrow \begin{cases} \tanh \tau \rightarrow 1 \\ \frac{1}{\cosh \tau} \rightarrow 0 \end{cases} \Rightarrow \tau \rightarrow \infty \Rightarrow \begin{cases} p \rightarrow \infty \\ \tau \rightarrow \infty \end{cases} \text{ or } \sigma \rightarrow 0.$$

$$3. c_{t/2} \rightarrow 0 \Rightarrow \frac{\sigma}{\cosh \tau} \rightarrow 0 \Rightarrow \begin{cases} \sigma \rightarrow 0 \\ \cosh \tau \rightarrow \infty \end{cases} \Rightarrow \begin{cases} p \rightarrow \infty \\ \tau \rightarrow \infty \end{cases} \text{ or } \sigma \rightarrow 0.$$

$$4. c_{t/2} \rightarrow \infty \Rightarrow \frac{\sigma}{\cosh \tau} \rightarrow \infty \Rightarrow \sigma \rightarrow \infty \Rightarrow x \rightarrow \bar{x} = 0.$$

$$5. t \rightarrow 0 \Rightarrow \frac{p}{\sigma} \rightarrow 0 \Rightarrow \begin{cases} p \rightarrow 0 \\ \sigma \rightarrow \infty \end{cases} \Rightarrow \begin{cases} \sigma \rightarrow 0 \\ x \rightarrow 0 \end{cases} \Rightarrow \sigma \rightarrow 0.$$

$$6. t \rightarrow \infty \Rightarrow \frac{p}{\sigma} \rightarrow \infty \Rightarrow \begin{cases} p \rightarrow \infty \\ \sigma \rightarrow 0 \end{cases} \Rightarrow \begin{cases} p \rightarrow \infty \\ \tau \rightarrow \infty \end{cases} \text{ or } \sigma \rightarrow 0.$$

$$7. |\alpha| \rightarrow \infty \Rightarrow \begin{cases} p \rightarrow \infty, \\ \tau \rightarrow \infty, \end{cases} \text{ or } x \rightarrow \bar{x} = 0 \quad \square$$

Lemma 3.10. *Suppose $\nu_n \in D_1 \cap N_3$. If $p \rightarrow \infty, \tau \rightarrow \infty, \sigma \rightarrow \bar{\sigma} \neq 0$, then $y \rightarrow \infty$.*

Proof. Since $\frac{2 \sinh(2p)}{\cosh(2p) + \cosh(2\tau)} < \infty$ for $p \rightarrow \infty$, then $y \rightarrow \infty$. \square

Lemma 3.11. *Suppose $\nu_n \in D_1 \cap N_3$. If $\sigma \rightarrow 0$, then one of the functions x, y, z, v tends to ∞ , otherwise x or z tends to 0.*

Proof. Assume the converse. We have $z \approx \frac{\cosh \tau (p \cosh p - \sinh p)}{\sigma^2 (\cosh(2p) + \cosh(2\tau))} \Rightarrow \begin{cases} p \rightarrow \infty, \\ p \rightarrow 0, \\ \tau \rightarrow \infty. \end{cases}$ So the proof is in these

three cases as follows:

1. $p \rightarrow \infty$. Then we get $y \approx \frac{1}{\sigma} p \rightarrow \infty$.

2. $p \rightarrow 0$. Consider three subcases:

2.1. $\tau \rightarrow 0$. Here we have $x \approx \frac{\tau p}{\sigma} \Rightarrow \sigma \approx \tau p$ and $y \approx \frac{p}{\sigma} \approx \frac{p}{\tau p} = \frac{1}{\tau} \rightarrow \infty$.

2.2. $\tau \rightarrow \infty$. We obtain

$$\begin{aligned} x &\approx \frac{p}{\sigma e^\tau} \Rightarrow p \approx \sigma e^\tau, \\ z &\approx \frac{p^3}{\sigma^2 e^\tau} \approx \frac{p^2}{\sigma} \Rightarrow \sigma \approx p^2, \\ y &\approx \frac{1}{\sigma} \left(\frac{p}{\cosh^2 \tau} - p \right) \approx \frac{p}{\sigma} \approx \frac{1}{p} \rightarrow \infty. \end{aligned}$$

2.3. $\tau \rightarrow \bar{\tau} < \infty$, $\bar{\tau} \neq 0$. It follows that $x \approx \frac{p}{\sigma} \Rightarrow p \approx \sigma$, then $z \approx \frac{p^3}{\sigma^2} \approx \sigma \rightarrow 0$.

3. $\tau \rightarrow \infty$, $p \rightarrow \bar{p} < \infty$, $\bar{p} \neq 0$. We get $\frac{\sinh(2p)}{\cosh(2p) + \cosh(2\tau)} \rightarrow 0 \Rightarrow y \approx \frac{\bar{p}}{\sigma} \rightarrow \infty$. □

Below in the case $\nu_n \in (D_1 \cap N_1) \cup (D_1 \cap N_2)$ we use the following notation:

$$s_i = \sin u_i, \quad c_i = \cos u_i, \quad d_i = \sqrt{1 - k^2 s_i^2}, \quad i = 1, 2. \quad (51)$$

$$E_1 = E(u_1, k), \quad F_1 = F(u_1, k), \quad \Delta = 1 - k^2 s_1^2 s_2^2. \quad (52)$$

Lemma 3.12. *Suppose $\nu_n \in (D_1 \cap N_1) \cup (D_1 \cap N_2)$. If $\Delta \rightarrow 0$, then $\frac{d_1 d_2}{\Delta}$ and $\frac{c_1 d_1}{\Delta}$ are bounded from above.*

Proof. Let $k^2 = 1 - c_3^2$; then $c_i \rightarrow 0$, $i = 1, 2, 3$. Introduce spherical coordinates as follows:

$$c_1 = r \sin \varphi_1 \cos \varphi_2, \quad c_2 = r \sin \varphi_1 \sin \varphi_2, \quad c_3 = r \cos \varphi_1.$$

Then it follows from $r \rightarrow 0$ that:

$$\begin{aligned} d_i^2 &= 1 - (1 - c_3^2)(1 - c_i^2) = c_i^2 + c_3^2 - c_i^2 c_3^2 \approx c_i^2 + c_3^2, \quad i = 1, 2, \\ \Delta &= 1 - (1 - c_3^2)(1 - c_1^2)(1 - c_2^2) \approx c_1^2 + c_2^2 + c_3^2 = r^2. \end{aligned}$$

This implies that

$$\begin{aligned} \left(\frac{d_1 d_2}{\Delta} \right)^2 &\approx \frac{(c_1^2 + c_3^2)(c_2^2 + c_3^2)}{r^4} = (\sin^2 \varphi_1 \cos^2 \varphi_2 + \cos^2 \varphi_1)(\sin^2 \varphi_1 \sin^2 \varphi_2 + \cos^2 \varphi_1) \leq 1, \\ \left(\frac{c_1 d_1}{\Delta} \right) &\approx \sin^2 \varphi_1 \cos^2 \varphi_2 (\sin^2 \varphi_1 \cos^2 \varphi_2 + \cos^2 \varphi_1) \leq 1. \end{aligned}$$

Therefore $\frac{d_1 d_2}{\Delta}$, $\frac{c_1 d_1}{\Delta}$ are bounded from above. □

In the next lemmas we use the following parametrization of exponential mapping for the case $\lambda \in C_1$ (see (15)):

$$\begin{aligned} x &= -\frac{4\sigma k s_1 s_2 d_1 d_2}{\alpha \Delta}, \\ y &= -\frac{4\sigma}{\alpha} \left(\frac{k^2 s_1 s_2^2 c_1 d_1}{\Delta} + \frac{F_1}{2} - E_1 \right), \\ z &= \frac{4k c_2 f_z}{|\alpha| \Delta}, & f_z &= c_1 (F_1 - 2E_1) + s_1 d_1, \\ v &= \frac{y^3}{6} - \frac{2k^2 (c_1 c_2 + s_1 s_2 d_1 d_2) y}{|\alpha| \Delta^2} + \frac{4}{3\alpha\sigma} \left(F_1 (1 - k^2) - E_1 (1 - 2k^2) \right. \\ &\quad \left. - \frac{k^2 s_1 d_1}{\Delta^3} \left(6s_1 s_2 c_2 d_1 d_2 (2d_2 - \Delta) + c_1 (1 + 3c_2^2 (d_2^2 - s_2^2) - k^4 s_1^2 s_2^6 (2d_1^2 + s_1^2)) \right) \right). \end{aligned}$$

Lemma 3.13. *If $\{\nu_n\} \subset D_1 \cap N_1$ satisfies $\nu_n \rightarrow \bar{\nu} \in \text{cl}(D_1) \setminus D_1$ and $\text{Exp}(\nu_n) \rightarrow \bar{q} \in M_1$, then $\Delta \rightarrow 0$ or $\sigma \rightarrow 0$.*

Proof. Notice that for $\bar{\nu} = (\bar{x}, \bar{y}, \bar{z}, \bar{v}) \in M_1$ we have $\bar{x} \neq 0$ and $\bar{z} \neq 0$. Consider all possible cases $\nu_n \rightarrow \partial D_1$:

1. $\theta_{t/2} \rightarrow 0 \Rightarrow \begin{cases} \sin \frac{\theta_{t/2}}{2} \rightarrow 0 \\ \cos \frac{\theta_{t/2}}{2} \rightarrow 1 \end{cases} \Rightarrow \begin{cases} k s_2 \rightarrow 0, \\ d_2 \rightarrow 1. \end{cases}$ It follows that $x \approx \frac{(k s_2) s_1 d_1}{\sigma} \Rightarrow \begin{cases} \sigma \rightarrow 0, \\ x \rightarrow \bar{x} = 0. \end{cases}$
2. $\theta_{t/2} \rightarrow \pi \Rightarrow \begin{cases} \sin \frac{\theta_{t/2}}{2} \rightarrow 1 \\ \cos \frac{\theta_{t/2}}{2} \rightarrow 0 \end{cases} \Rightarrow \begin{cases} k s_2 \rightarrow 1, \\ d_2 \rightarrow 0. \end{cases}$ Then $x \approx \frac{s_1 d_1 d_2}{\sigma \Delta} \Rightarrow \begin{cases} \sigma \Delta \rightarrow 0, \\ x \rightarrow \bar{x} = 0. \end{cases}$
3. $c_{t/2} \rightarrow 0 \Rightarrow k \sigma c_2 \rightarrow 0 \Rightarrow z \approx \frac{k c_2 f_z}{\sigma^2 \Delta} \Rightarrow \frac{\sigma^3 \Delta}{f_z} \rightarrow 0$, otherwise $z \rightarrow \bar{z} = 0$. This means that $\sigma^3 \Delta \rightarrow 0$ or $f_z \rightarrow \infty$. Suppose $f_z \rightarrow \infty$, $\sigma \rightarrow \bar{\sigma} \neq 0$, then $u_1 \rightarrow \pi/2$ and $k \rightarrow 1$. Since $k \sigma c_2 \rightarrow 0$, then $u_2 \rightarrow \pi/2 \Rightarrow \Delta \rightarrow 0$.
4. $c_{t/2} \rightarrow \infty \Rightarrow k \sigma c_2 \rightarrow \infty \Rightarrow \sigma \rightarrow \infty \Rightarrow \Delta \rightarrow 0$, otherwise $x \rightarrow \bar{x} = 0$.
5. $t \rightarrow 0 \Rightarrow \frac{p}{\sigma} \rightarrow 0 \Rightarrow \begin{cases} u_1 \rightarrow 0 \\ \sigma \rightarrow \infty \end{cases} \Rightarrow \Delta \rightarrow 0$, otherwise $x \rightarrow \bar{x} = 0$.
6. $t \rightarrow t_{\text{MAX}}^1 \Rightarrow \begin{cases} f_z(u_1, k) \rightarrow 0 \text{ for } k \geq k_0 \\ u_1 \rightarrow \pi \text{ for } k \leq k_0 \end{cases} \Rightarrow \begin{cases} \sigma^2 \Delta \rightarrow 0, \text{ otherwise } z \rightarrow \bar{z} = 0 \\ \sigma \Delta \rightarrow 0, \text{ otherwise } x \rightarrow \bar{x} = 0. \end{cases}$
7. $|\alpha| \rightarrow \infty \Rightarrow \sigma \rightarrow \infty \Rightarrow \Delta \rightarrow 0$, otherwise $x \rightarrow \bar{x} = 0$. □

Lemma 3.14. *Suppose $\nu_n \in D_1 \cap N_1$. If $\Delta \rightarrow 0$, then $x \rightarrow 0$ or $y \rightarrow \infty$.*

Proof. Consider two possible cases:

1. $\sigma \rightarrow \infty \Rightarrow x = -4k s_1 s_2 \frac{d_1 d_2}{\Delta} \frac{1}{\sigma} \rightarrow 0$ (see Lemma 3.12).
2. $\sigma \rightarrow \bar{\sigma} < \infty$. It follows from Lemma 3.12 that $k^2 s_1 s_2^2 \frac{c_1 d_1}{\Delta} - E_1$ is bounded from above. Since $F_1 \rightarrow \infty$, then $y \rightarrow \infty$. □

Lemma 3.15. *Suppose $\nu_n \in D_1 \cap N_1$. If $\sigma \rightarrow 0, \Delta \rightarrow \bar{\Delta} \neq 0$, then one of the functions x, y, z, v tends to ∞ , otherwise x or z tends to zero.*

Proof. Assume the converse. Then notice that $k s_1 s_2 d_1 d_2 \rightarrow 0$, otherwise $x \approx \frac{1}{\sigma} \rightarrow \infty$. The proof consists of the following six items:

1. $d_1 \rightarrow 0$. This means that $u_1 \rightarrow \pi/2, k \rightarrow 1$ and $u_2 \rightarrow \bar{u}_2 \neq \pi/2$. Whence, $y \approx F_1/\sigma \rightarrow \infty$.

2. $u_1 \rightarrow 0$. Consider four subcases:

2.1. $s_2 k \rightarrow 0$. Here we have

$$x \approx \frac{s_2 k u_1}{\sigma} \Rightarrow s_2 k u_1 \approx \sigma,$$

$$F_1 \sim u_1, E_1 \sim u_1 \Rightarrow y \approx \frac{1}{\sigma} \left(s_1 (s_2 k)^2 \frac{c_1 d_1}{\Delta} + \frac{F_1}{2} - E_1 \right) \approx \frac{u_1}{\sigma} \approx \frac{1}{s_2 k} \rightarrow \infty.$$

2.2. $d_2 \rightarrow 0$. It follows that

$$x \approx \frac{d_2 u_1}{\sigma} \Rightarrow d_2 u_1 \approx \sigma, \quad y \approx \frac{u_1}{\sigma} \approx \frac{1}{d_2} \rightarrow \infty.$$

2.3. $c_2 \rightarrow 0$, $k \rightarrow \bar{k} \in (0, 1)$. We get

$$x \approx \frac{u_1}{\sigma} \Rightarrow u_1 \approx \sigma, \quad z \approx \frac{c_2 u_1^3}{\sigma^2} \approx c_2 u_1 \rightarrow 0.$$

2.4. $k \rightarrow \bar{k} \neq 0$, $u_2 \rightarrow \bar{u}_2 \in (0, \pi/2)$. Hence

$$x \approx \frac{u_1}{\sigma}, \quad z \approx \frac{u_1^3}{\sigma^2} \approx x^2 u_1 \rightarrow 0.$$

3. $u_1 \rightarrow \pi \Rightarrow F_1 \rightarrow 2K$, $E_1 \rightarrow 2E$. We obtain $y \approx \frac{2E - K}{\sigma} \Rightarrow 2E - K \rightarrow 0$. It follows that $v \approx \frac{K}{\sigma^3} \rightarrow \infty$, since $K \rightarrow \bar{K} = K(\bar{k}) > 0$.

4. $d_2 \rightarrow 0$, $u_1 \rightarrow \bar{u}_1 \in (0, \pi/2) \Rightarrow u_2 \rightarrow \pi/2$, $k \rightarrow 1$.

$$x \approx \frac{d_2}{\sigma}, \quad y \approx \frac{1}{\sigma} \left(s_1 + \frac{F_1}{2} - E_1 \right).$$

Note that the function $\sin u_1 + \frac{F(u_1, 1)}{2} - E(u_1, 1)$ vanishes only at the point $u_1 = 0$ since it has positive derivative $\frac{1}{2\sqrt{1 - \sin^2 u_1}}$. Therefore $y \rightarrow \infty$.

5. $u_2 \rightarrow 0$, $s_1 \rightarrow \bar{s}_1 \neq 0$, $d_1 \rightarrow \bar{d}_1 \neq 0$, $k \rightarrow \bar{k} \neq 0$. We have $y \approx \frac{2E_1 - F_1}{\sigma} \Rightarrow 2E_1 - F_1 \rightarrow 0$. Hence $z \approx \frac{1}{\sigma^2} \rightarrow \infty$.

6. $k \rightarrow 0$. We get $y \approx \frac{u_1}{\sigma} \Rightarrow u_1 \rightarrow 0$ (see item 2.1). □

In the next lemmas we use the following parametrization of exponential mapping for the case $\lambda \in C_2$ (see (16)):

$$\begin{aligned}
x &= -\frac{4 \operatorname{sgn} c \sigma s_1 s_2 c_1 c_2}{\alpha k \Delta}, \\
y &= -\frac{4\sigma}{\alpha k} \left(\frac{k^2 s_1 s_2^2 c_1 d_1}{\Delta} + \left(1 - \frac{k^2}{2}\right) F_1 - E_1 \right), \\
z &= -4 \frac{4 \operatorname{sgn} c d_2 g_z}{|\alpha| k^2 \Delta}, \quad g_z = (2E_1 + (k^2 - 2)F_1)d_1 - k^2 s_1 c_1, \\
v &= \frac{y^3}{6} + \frac{2y}{|\alpha| k^2 \Delta^2} \left(1 + (1 + c_1^2 c_2^2) k^4 s_1^2 s_2^2 - k^2 (s_1^2 + s_2^2 - 2c_1 c_2 d_1 d_2 s_1 s_2) \right) \\
&\quad - \frac{3}{4\alpha\sigma k} \left(2F_1 \left(\frac{1}{k^2} - 1 \right) - E_1 \left(\frac{2}{k^2} - 1 \right) + \frac{c_1 s_1}{\Delta^3} \left(2c_2^2 d_1 (1 + d_1^2) \Delta + 6c_1 c_2 d_2 k^2 s_1 s_2 (2c_2^2 - \Delta) \right. \right. \\
&\quad \left. \left. + d_1 s_2^2 ((2 - k^2) \Delta^2 - 4d_1^2 d_2^2) \right) \right).
\end{aligned}$$

Lemma 3.16. *If $\{\nu_n\} \subset D_1 \cap N_2$ satisfies $\nu_n \rightarrow \bar{\nu} \in \operatorname{cl}(D_1) \setminus D_1$ and $\operatorname{Exp}(\nu_n) \rightarrow \bar{q} \in M_1$, then $\Delta \rightarrow 0$ or $\frac{\sigma}{k} \rightarrow 0$.*

Proof. Notice that for $\bar{\nu} = (\bar{x}, \bar{y}, \bar{z}, \bar{v}) \in M_1$ we have $\bar{x} \neq 0$. Consider all possible cases for $\bar{\nu} \in \operatorname{cl}(D_1) \setminus D_1$:

1. $\theta_{t/2} \rightarrow 0 \Rightarrow \begin{cases} \sin \frac{\theta_{t/2}}{2} \rightarrow 0 \\ \cos \frac{\theta_{t/2}}{2} \rightarrow 1 \end{cases} \Rightarrow \begin{cases} s_{u_2} \rightarrow 0, \\ c_{u_2} \rightarrow 1. \end{cases}$ It follows that $\frac{\sigma\Delta}{k} \rightarrow 0$, otherwise $x \rightarrow \bar{x} = 0$.
2. $\theta_{t/2} \rightarrow \pi \Rightarrow \begin{cases} \sin \frac{\theta_{t/2}}{2} \rightarrow 1 \\ \cos \frac{\theta_{t/2}}{2} \rightarrow 0 \end{cases} \Rightarrow \begin{cases} s_2 \rightarrow 1, \\ c_2 \rightarrow 0. \end{cases}$ This means that $\frac{\sigma\Delta}{k} \rightarrow 0$, otherwise $x \rightarrow \bar{x} = 0$.
3. $c_{t/2} \rightarrow 0 \Rightarrow \frac{\sigma}{k} d_2 \rightarrow 0 \Rightarrow \frac{\sigma}{k} \rightarrow 0$ or $c_2 \rightarrow 0$. From $c_2 \rightarrow 0$ we have $\frac{\sigma\Delta}{k} \rightarrow 0$, otherwise $x \rightarrow \bar{x} = 0$.
4. $c_{t/2} \rightarrow \infty \Rightarrow \frac{\sigma}{k} d_2 \rightarrow \infty \Rightarrow \frac{\sigma}{k} \rightarrow \infty \Rightarrow \Delta \rightarrow 0$, otherwise $x \rightarrow \bar{x} = 0$.
5. $t \rightarrow 0 \Rightarrow \frac{pk}{\sigma} \rightarrow 0 \Rightarrow \frac{s_1 k}{\sigma} \rightarrow 0 \Rightarrow \Delta \rightarrow 0$, otherwise $x \rightarrow \bar{x} = 0$.
6. $t \rightarrow t_{\text{MAX}}^1 \Rightarrow u_1 \rightarrow \frac{\pi}{2} \Rightarrow \frac{\sigma\Delta}{k} \rightarrow 0$, otherwise $x \rightarrow \bar{x} = 0$.
7. $\alpha \rightarrow \infty \Rightarrow \sigma \rightarrow \infty \Rightarrow \Delta \rightarrow 0$, otherwise $x \rightarrow \bar{x} = 0$. □

Lemma 3.17. *Suppose $\nu_n \in D_1 \cap N_2$. If $\Delta \rightarrow 0$, then $x \rightarrow 0$ or $y \rightarrow \infty$.*

Proof. Consider two possible cases:

1. $\sigma \rightarrow \infty \Rightarrow x = -4k s_1 s_2 \frac{c_1 c_2}{\Delta} \frac{1}{\sigma} \rightarrow 0$ (see Lemma 3.12).
2. $\sigma \rightarrow \bar{\sigma} < \infty$. It follows from Lemma 3.12 that $(k^2 s_1 s_2^2 \frac{c_1 d_1}{\Delta} - E_1)$ is bounded from above. And since $F_1 \rightarrow \infty$ we have $y \rightarrow \infty$. □

Lemma 3.18. *Suppose $\nu_n \in D_1 \cap N_2$. If $\frac{\sigma}{k} \rightarrow 0, \Delta \rightarrow \bar{\Delta} \neq 0$, then one of the functions x, y, z or v tends to ∞ , otherwise x or z tends to 0.*

Proof. Assume the converse. Then notice that $s_1 s_2 c_1 c_2 \rightarrow 0$, otherwise $x \approx \frac{k}{\sigma} \rightarrow \infty$. The proof consists of five steps:

1. $u_1 \rightarrow 0$. Then we obtain

$$x \approx \frac{u_1 s_1 c_2}{\sigma/k} \Rightarrow \frac{\sigma}{k} \approx u_1 s_2 c_2.$$

It follows from Taylor expansion that $g_z \approx k^2 u_1^3$, then

$$z \approx \frac{d_2 k^2 u_1^3}{\sigma^2} \approx \frac{d_2 u_1}{s_2^2 c_2^2} = \frac{d_2 u_1}{s_2^2 c_2^2} \Rightarrow s_2 c_2 \rightarrow 0,$$

otherwise $z \rightarrow 0$.

$$y \approx \frac{1}{\sigma k} \left(k^2 u_1 s_2^2 + (1 - k^2/2)u_1 - u_1 \right) = \frac{u_1 (s_2^2 - 1/2)}{\sigma/k} \approx \frac{s_2^2 - 1/2}{s_2 c_2} \rightarrow \infty.$$

2. $k \rightarrow 0$, $u_1 \rightarrow \bar{u}_1 \neq 0$. Using Taylor expansion we get $z \approx \frac{k^2}{\sigma^2} \rightarrow \infty$.

3. $u_1 \rightarrow \frac{\pi}{2}$, $k \rightarrow \bar{k} \neq 0$. We have $y \approx \frac{1}{\sigma} \left(\frac{k^2 c_1 s_1 d_1^2 s_2^2}{\Delta} + (1 - k^2/2)F_1 - E_1 \right)$. Notice that

$$\frac{d}{dk} \left((1 - k^2/2)F_1 - E_1 \right) = \frac{k}{2(1 - k^2)} \int_0^{\pi/2} \frac{k^2 \cos^2 \theta d\theta}{\sqrt{1 - k^2 \sin^2 \theta}} > 0 \Rightarrow (1 - k^2/2)F_1 - E_1 > 0.$$

Combining the last inequality and $\frac{k^2 c_1 s_1 d_1^2 s_2^2}{\Delta} \rightarrow 0$, we obtain $y \rightarrow \infty$.

4. $u_2 \rightarrow 0$, $u_1 \rightarrow \bar{u}_1 \in (0, \pi/2)$, $k \rightarrow \bar{k} \neq 0$. Here we have

$$x \approx \frac{u_2}{\sigma} \Rightarrow u_2 \approx \sigma,$$

otherwise $x \rightarrow 0$.

$$z \approx \frac{(2E_1 + (k^2 - 2)F_1)d_1 - k^2 c_1 s_1}{\sigma^2},$$

$$y \approx \frac{1}{\sigma} \left(k^2 s_1 c_1 d_1 s_2^2 + (1 - k^2/2)F_1 - E_1 \right).$$

Since $k^2 s_1 c_1 d_1 s_2^2 \rightarrow 0$, we see that $(1 - k^2/2)F_1 - E_1 \rightarrow 0$, otherwise $y \rightarrow \infty$. Hence from $k^2 c_1 s_1 \approx 1$ we get $z \approx \frac{1}{\sigma^2} \rightarrow \infty$.

5. $u_2 \rightarrow \pi/2$, $u_1 \rightarrow u_1 \in (0, \pi/2)$, $k \rightarrow \bar{k} \neq 0$. Suppose $\bar{k} \neq 1$, then $z \approx \frac{d_2 g_z}{\sigma^2} \rightarrow \infty$. This means that $\bar{k} = 1$. Here we have $y \approx \frac{1}{\sigma} \left(\frac{c_1 s_1 d_1}{\Delta} + F_1/2 - E_1 \right) = \frac{1}{\sigma} (s_1 + F_1/2 - E_1)$. Since $\frac{d}{du_1} (s_1 + F_1/2 - E_1) = \frac{1}{2 \cos u_1} > 0$, it follows that $y \rightarrow \infty$. \square

Theorem 3.19. *The mapping $\text{Exp} : D_i \rightarrow M_i$ is proper for $i = 1, \dots, 4$.*

Proof. Assume the converse. Then it follows from Lemma 3.1 that $\text{Exp} : D_1 \rightarrow M_1$ is not proper. By Lemma 3.2, there exists a sequence $\nu_n \in D_1$, such that $\nu_n \rightarrow \bar{\nu} \in \text{cl}(D_1) \setminus D_1$, $\text{Exp}(\nu_n) \rightarrow \bar{q} \in M_1$. Since $\nu_n \in D_1$, we consider 3 cases:

- (1) $\{\nu_n\} \subset D_1 \cap N_6$ is impossible (see Lemmas 3.7, 3.8),
- (2) $\{\nu_n\} \subset D_1 \cap N_3$ is impossible (see Lemmas 3.9–3.11),
- (3) $\{\nu_n\} \subset (D_1 \cap N_1) \cup (D_1 \cap N_2)$ is impossible (see Lemmas 3.13–3.18).

Since all cases are impossible, we have a contradiction which proves the theorem. \square

Theorem 3.20. *The mapping $\text{Exp} : D_i \rightarrow M_i$ is a diffeomorphism for $i = 1, \dots, 4$.*

Proof. Follows from Th. 2.5, since all hypotheses of this theorem hold by Propos. 2.7, Th. 1.3 and Th. 3.19. \square

Corollary 3.21. *The mapping $\text{Exp} : \tilde{N} \rightarrow \tilde{M}$ is a diffeomorphism.*

4. CUT TIME

In this section we prove that the cut time coincides with the first Maxwell time corresponding to reflections.

4.1. Cut time and Maxwell time

Theorem 4.1. *For any $\lambda \in C$,*

$$t_{\text{cut}}(\lambda) = t_{\text{MAX}}^1(\lambda).$$

Proof. Take any $\lambda \in C$ and denote $t_1 = t_{\text{MAX}}^1(\lambda)$. Since $t_{\text{cut}}(\lambda) \leq t_1$ by Th. 1.1, it remains to prove that $t_{\text{cut}}(\lambda) \geq t_1$.

Let us call a pair $(\lambda, t) \in N$ optimal if the geodesic $\text{Exp}(\lambda, s)$ is optimal on the segment $s \in [0, t]$. We have to show that (λ, t) is optimal for any $t \in (0, t_1)$.

- (1) If $\lambda \in C_4 \cup C_5 \cup C_7$, then $t_1 = +\infty$, and any (λ, t) , $t \in (0, t_1)$, is optimal since (x_s, y_s) is a straight line.
- (2) Let $\lambda \in C_1 \cup C_2 \cup C_6$, thus $t_1 \in (0, +\infty)$.

Since $t_1 = t_{\text{MAX}}^1(\lambda)$, then $\nu_1 = (\lambda, t_1) \in N'$. For $\lambda \in C_1 \cup C_2 \cup C_6$ the function $t \mapsto \sin \theta_{t/2} c_{t/2}$ has isolated zeros, thus there exists $t \in (0, t_1)$ arbitrarily close to t_1 such that $\nu = (\lambda, t) \in \tilde{N}$. Then $q = \text{Exp}(\nu) \in \tilde{M}$ (Propos. 2.4). Since $\text{Exp}(N') \cap \tilde{M} = \emptyset$ (Propos. 2.3) and $\text{Exp} : \tilde{N} \rightarrow \tilde{M}$ is a diffeomorphism (see Corollary 3.21) then $\text{Exp}^{-1}(q) \cap \tilde{N} = \{\nu\}$. Thus $\nu = (\lambda, t)$ is optimal. Since t can be chosen arbitrarily close to t_1 , then any (λ, t) , $t \in (0, t_1)$, is optimal.

- (3) Let $\lambda \in C_3$, then $t_1 = +\infty$. There exist $(\lambda, t) \in \tilde{N}$ for arbitrarily large t . Then the proof follows the argument of item (2).

\square

Now we collect all properties of the cut time that we previously obtained for the Maxwell time t_{MAX}^1 .

Corollary 4.2. *The function $t_{\text{cut}} : C \rightarrow (0, +\infty]$ has the following properties:*

- (1) *Let $\lambda \in C$ and let $t_1 = t_{\text{cut}}(\lambda)$. For finite t_1 , a trajectory $\text{Exp}(\lambda, s)$, $s \in [0, t]$, is optimal iff $t \in [0, t_1]$. For $t_1 = +\infty$, any trajectory $\text{Exp}(\lambda, s)$, $s \in [0, t]$, $t > 0$, is optimal.*

(2) The function t_{cut} has the following explicit representation:

$$\begin{aligned} \forall \lambda \in C_1 & & t_{\text{cut}}(\lambda) &= \frac{p_1(k)}{\sqrt{|\alpha|}}, \\ \forall \lambda \in C_2 & & t_{\text{cut}}(\lambda) &= \frac{2Kk}{\sqrt{|\alpha|}}, \\ \forall \lambda \in C_6 & & t_{\text{cut}}(\lambda) &= \frac{2\pi}{\sqrt{|c|}}, \\ \forall \lambda \in C_3 \cup C_4 \cup C_5 \cup C_7 & & t_{\text{cut}}(\lambda) &= +\infty. \end{aligned}$$

(3) The function t_{cut} depends only on E and $|\alpha|$, is preserved by the flow of \vec{H}_v and by the reflections ε^i , and is homogeneous of order one w.r.t. the dilations δ_μ .

(4) The function t_{cut} is continuous on $C \setminus C_4$ and is smooth on $C_1^0 \cup C_2$, where $C_1^0 = \{\lambda \in C_1 \mid k \neq k_0\}$.

According to Cor. 4.2, the function $t_{\text{cut}} : C \rightarrow (0, +\infty]$ is invariant w.r.t. the flow $e^{s\vec{H}_v}$ and the reflections ε^i , and respects the action of dilations δ_μ :

$$\begin{aligned} t_{\text{cut}} \circ e^{s\vec{H}_v} &= t_{\text{cut}} \circ \varepsilon^i = t_{\text{cut}}, \\ t_{\text{cut}} \circ \delta_\mu &= \mu t_{\text{cut}}. \end{aligned}$$

Thus the cut time can be represented (up to a constant positive factor) by a univariate function on the quotient $C / \langle e^{s\vec{H}_v}, \varepsilon^i, \delta_\mu \rangle$. The quotient $C / \langle e^{s\vec{H}_v}, \varepsilon^i \rangle$ can be represented by the quadrant $\{(\theta, c, \alpha) \in C \mid \theta = 0, c \geq 0, \alpha \geq 0\}$, thus

$$C / \langle e^{s\vec{H}_v}, \varepsilon^i, \delta_\mu \rangle \cong \Gamma \sqcup P,$$

where

$$\begin{aligned} \Gamma &= \{(\theta, c, \alpha) \in C \mid \theta = 0, c = \sin \beta, \alpha = \cos \beta, \beta \in [0, \pi/2]\}, \\ P &= \{(\theta, c, \alpha) \in C \mid \theta = 0, c = 0, \alpha = 0\}. \end{aligned}$$

The point P corresponds to the subset C_7 , while the arc Γ corresponds to the rest subsets $C \setminus C_7 = C_1 \cup C_2 \cup C_{35} \cup C_4 \cup C_6$ of decomposition (36).

Thus (up to a constant positive factor) the cut time can be represented on the set $C \setminus C_7$ as a univariate function $t_{\text{cut}}(\beta)$, $\beta \in [0, \pi/2]$.

If $\beta = 0$, then $\lambda \in C_4$, thus $t_{\text{cut}}(\beta) = +\infty$.

If $\beta \in (0, \beta_1)$, where $\beta_1 = \arccos(\sqrt{5} - 2)$, then $\lambda \in C_1$, thus

$$t_{\text{cut}}(\beta) = \frac{2p_1(k)}{\sqrt{\alpha}}, \quad k = \sqrt{\frac{\sin^2 \beta}{4 \cos \beta}}, \quad \alpha = \cos \beta, \quad p_1(k) = \min(p_z^1(k), 2K(k)).$$


If $\beta = \beta_1$, i.e., $\sin^2 \beta = 4 \cos \beta$, then $\lambda \in C_{35}$, thus $t_{\text{cut}}(\beta) = +\infty$.

If $\beta \in (\beta_1, \pi/2)$, then $\lambda \in C_2$, thus

$$t_{\text{cut}}(\beta) = \frac{2Kk}{\sqrt{\alpha}}, \quad k = \sqrt{\frac{4 \cos \beta}{\sin^2 \beta}}, \quad \alpha = \cos \beta.$$

Finally, if $\beta = \pi/2$, then $\lambda \in C_6$, thus $t_{\text{cut}}(\beta) = 2\pi$.

The plot of the function $t_{\text{cut}}(\beta)$ is shown in Fig. 3. Notice continuity of $t_{\text{cut}}(\beta)$ everywhere except $\beta = 0$,

FIGURE 3. Plot of the function $\beta \mapsto t_{\text{cut}}(\beta)$

where $t_{\text{cut}}(+0) = 2\pi < +\infty = t_{\text{cut}}(0)$. Also notice smoothness of $t_{\text{cut}}(\beta)$ everywhere except $\beta = 0$, $\beta = \beta_1$, and $\beta = \beta_0$, where $\frac{\sin^2 \beta_0}{4 \cos \beta_0} = k_0^2$, $2E(k_0) - K(k_0) = 0$ (here $E(k_0)$ is the complete elliptic integral of the second kind), corresponds to the figure-of-eight closed Euler elastica. These regularity properties of $t_{\text{cut}}(\lambda) = t_{\text{MAX}}^1(\lambda)$ are reported in Corollary 4.2.

4.2. Cut time and conjugate time

Proposition 4.3. *Let $\lambda \in C_1$, $t_1 = t_{\text{cut}}(\lambda)$, $\tau = (\varphi + t_1/2)/\sqrt{|\alpha|}$. Then $t_1 = t_{\text{conj}}^1(\lambda)$ iff one of the conditions hold:*

- (1) $k < k_0$, $\text{sn } \tau = 0$,
- (2) $k = k_0$,
- (3) $k > k_0$, $\text{cn } \tau = 0$.

In particular, if $t_1 = t_{\text{conj}}^1(\lambda)$, then $\text{sn } \tau \text{ cn } \tau = 0$ or $k = k_0$.

Proof. Follows immediately from Lemma 8 [2]. □

Remark 4.4. The equality $\text{sn } \tau = 0$ ($\text{cn } \tau = 0$) is equivalent to $\sin \theta_{t/2} = 0$ ($c_{t/2} = 0$); it means that elastica (x_t, y_t) , $t \in [0, t_1]$, is centered at a vertex (resp. inflexion point). The equality $k = k_0$ means that (x_t, y_t) , $t \in [0, t_1]$, is the closed figure-of-eight elastica.

Proposition 4.5. *Let $\lambda \in C_2$, $t_1 = t_{\text{cut}}(\lambda)$, $\tau = (\varphi + t_1/2)/(k\sqrt{|\alpha|})$. Then $t_1 = t_{\text{conj}}^1(\lambda)$ iff $\text{sn } \tau \text{ cn } \tau = 0$.*

Proof. Follows immediately from Lemma 8 [2]. □

Remark 4.6. The equality $\text{sn } \tau \text{ cn } \tau = 0$ is equivalent to $\sin \theta_{t/2} = 0$. It means that the corresponding elastica is centered at vertex.

Proposition 4.7. *Let $\lambda \in C_6$, $t_1 = t_{\text{cut}}(\lambda)$. Then $t_1 = t_{\text{conj}}^1(\lambda)$ iff $\sin \theta = 0$.*

Proof. Let $\lambda = (\theta, c, \alpha) \in C_6$, $\alpha = 0$, $c \neq 0$, $t_1 = t_{\text{cut}}(\lambda)$, $\nu = (\lambda, t_1) \in N_6$. Since $C_6 \subset \text{cl}(C_2)$, the expression for Jacobian $J = \frac{\partial(x, y, z, v)}{\partial(\theta, c, \alpha, t)}$ for $\nu \in N_6$ can be obtained by passing to the limit $\alpha \rightarrow 0$ in the expression for Jacobian $J|_{N_2}$ computed in [2]. By such a limit we get $J(\nu) = \frac{\pi^3}{|c|^3} \sin^2 \theta$. So the instant t_1 is a conjugate time iff $\sin \theta = 0$. \square

4.3. Optimal trajectories for special boundary conditions

For a generic terminal point $q_1 \in \widetilde{M}$, there exists a unique optimal trajectory $q_t = \text{Exp}(\lambda, t)$, $t \in [0, t_1]$, which can be found by solving the equation $\text{Exp}(\lambda, t_1) = q_1$, $(\lambda, t_1) \in \widetilde{N}$.

In this subsection we discuss special boundary conditions for which optimal trajectories can be given explicitly or by a more simple equation.

4.3.1. Abnormal variety

Consider the set of points in M filled by abnormal trajectories:

$$A = \{q \in M \mid x = z = 0, v = y^3/6\}.$$

We have $\text{Exp}(C_4, \mathbb{R}_+) = \text{Exp}(C_5, \mathbb{R}_+) = \text{Exp}(C_7^{0,\pi}, \mathbb{R}_+) = A \setminus \{q_0\}$, where

$$C_7^{0,\pi} = \{\lambda = (\theta, c, \alpha) \in C_7 \mid \alpha = c = 0, \theta \in \{0, \pi\}\}.$$

Any nonzero point $q_1 = (0, y_1, 0, v_1) \in A$ is connected with q_0 by a unique optimal trajectory

$$x_t = 0, \quad y_t = t \operatorname{sgn} y_1, \quad z_t = 0, \quad v_t = \frac{t^3}{6} \operatorname{sgn} y_1, \quad t \in [0, |y_1|].$$

4.3.2. Straight lines (x_t, y_t)

The set of points in M filled by trajectories that project to straight lines (x_t, y_t) is

$$L = \{q \in M \mid z = 0, v = (x^2 + y^2)y/6\} \supset A.$$

We have $\text{Exp}(C_7, \mathbb{R}_+) = L \setminus \{q_0\}$. The unique optimal trajectory for $q_1 \in L \setminus \{q_0\}$ is $\text{Exp}(\lambda, t)$, $\lambda \in C_7$, i.e.,

$$x_t = -t \sin \theta, \quad y_t = t \cos \theta, \quad z_t = 0, \quad v_t = \frac{t^3}{6} \cos \theta, \quad t \in [0, t_1],$$

where $t_1 > 0$ and $\theta \in S^1$ are found from the equations

$$x_1 = -t_1 \sin \theta, \quad y_1 = t_1 \cos \theta.$$

4.3.3. Fixed points of reflection ε^6

The reflection $\varepsilon^6: M \rightarrow M$ has the set of fixed points

$$S_6 = \{q \in M \mid y = 0, v = xz/2\}.$$

This 2-dimensional manifold is of particular interest since it is the only fixed manifold of reflections

$$S_i = \{q \in M \mid \varepsilon^i(q) = q\}, \quad i = 1, \dots, 7,$$

not contained completely in codimension one manifolds $S_1 = \{q \in M \mid z = 0\}$ and $S_2 = \{q \in M \mid x = 0\}$. By Lem. 4 [1],

$$S_3 \cup S_4 \cup S_5 \subset S_2, \quad S_7 \subset S_1.$$

But $S_6 \not\subset S_1 \cup S_2$.

If a point $q_1 \in S_6 \setminus \{q_0\}$ is connected with q_0 by a trajectory $q_t = \text{Exp}(\lambda, t)$, $t \in [0, t_1]$, $q_{t_1} = q_1$, then the trajectory $q_t^6 = \text{Exp}(\lambda^6, t)$, $t \in [0, t_1]$, satisfies the equation $q_{t_1}^6 = q_1$.

Moreover, if $\lambda^6 \neq \lambda$, then the points $(\lambda, t_1), (\lambda^6, t_1)$ would belong to the Maxwell set

$$\text{MAX}^6 = \{(\lambda, t) \in N \mid \lambda^i \neq \lambda, \text{Exp}(\lambda^i, t) = \text{Exp}(\lambda, t)\},$$

which might a priori give Maxwell times which are additional to those provided by the sets $\text{MAX}^1, \text{MAX}^2$ studied in [1]. It turns out that, as we show below, the equality $\lambda^6 = \lambda$ is satisfied for all $\lambda \in C$ with $\text{Exp}(\lambda, t_1) \in S_6$, $t_1 > 0$.

Consider the decomposition

$$S_6 = \bigsqcup_{i,j \in \{0,+,-\}} S_{ij},$$

$$S_{ij} = \{q \in S_6 \mid \text{sgn } x = i, \text{sgn } z = j\}.$$

For example, $S_{+-} = \{q \in S_6 \mid x > 0, z < 0\}$.

Denote $N_{ij} = \{(\lambda, t) \in N_6 \mid \tau = -i\pi/2, \text{sgn } c = j, t \in (0, 2\pi/|c|)\}$, $i, j \in \{+, -\}$, where $\tau = \theta + ct/2$.

Lemma 4.8. *For any $i, j \in \{+, -\}$, the mapping $\text{Exp}: N_{ij} \rightarrow S_{ij}$ is a diffeomorphism.*

Proof. The reflections ε^4 and ε^7 permute the sets N_{ij}, S_{ij} , thus it suffices to prove only that the mapping $\text{Exp}: N_{++} \rightarrow S_{++}$ is a diffeomorphism.

If $(\lambda, t) \in N_{++}$, then

$$x_t = \frac{2 \sin p}{c}, \quad y_t = 0, \quad z_t = \frac{2p - \sin(2p)}{2c^2}, \quad v_t = \frac{x_t z_t}{2}, \quad (53)$$

where $p = ct/2 \in (0, \pi)$. Thus $\text{Exp}(N_{++}) \subset S_{++}$.

Further, the mapping $\Phi: (p, c) \mapsto (x, z)$ is a diffeomorphism from $(0, \pi) \times \mathbb{R}_+$ to $\mathbb{R}_+ \times \mathbb{R}_+$ by Hadamard global diffeomorphism theorem, thus $\text{Exp}: N_{++} \rightarrow S_{++}$ is a diffeomorphism as well. \square

Denote $N^i_{0j} = \{(\lambda, t) \in N_6 \mid \tau = i\pi/2, \text{sgn } c = j, t = 2\pi/|c|\}$, $i, j \in \{+, -\}$.

Lemma 4.9. *Each of the mappings $\text{Exp}: N^+_{0j} \rightarrow S_{0j}$, $\text{Exp}: N^-_{0j} \rightarrow S_{0j}$, $j \in \{+, -\}$, is a diffeomorphism.*

Proof. Follows from the parameterization of trajectories (53) with $p = \pm\pi$. \square

Denote $N_{i0} = \{(\lambda, t) \in N_7 \mid \theta = -i\pi/2\}$, $i \in \{+, -\}$.

Lemma 4.10. *The mappings $\text{Exp}: N_{i0} \rightarrow S_{i0}$, $i \in \{+, -\}$, are diffeomorphisms.*

Proof. Follows immediately from the parameterization of extremal trajectories for $\lambda \in C_7$. \square

Lemmas 4.8–4.10 yield the following optimal synthesis for the terminal manifold S_6 .

Corollary 4.11. *Let $q_1 \in S_6 \setminus \{q_0\}$.*

(1) *If $q_1 \in S_{ij}$, $i, j \in \{+, -\}$, then the only optimal trajectory is $\text{Exp}(\lambda, t)$, $t \in [0, t_1]$, where $(\lambda, t_1) \in N_{ij}$ is determined by the equations*

$$x_1 = i \frac{\sin p}{c}, \quad z_1 = \frac{2p - \sin(2p)}{2c^2}, \quad jp \in (0, \pi), \quad jc \in (0, +\infty).$$

- (2) If $q_1 \in S_{0j}$, $j \in \{+, -\}$, then there are two optimal trajectories $\text{Exp}(\lambda_+, t)$, $\text{Exp}(\lambda_-, t)$, $t \in [0, t_1]$, where $(\lambda_{\pm}, t_1) \in N_{0j}^{\pm}$ is determined by the equations

$$z_1 = \frac{j\pi}{c^2}, \quad \tau = \pm \frac{\pi}{2}.$$

- (3) If $q_1 \in S_{i0}$, $i \in \{+, -\}$, then the only optimal trajectory is $\text{Exp}(\lambda, t)$, $t \in [0, t_1]$, where $(\lambda, t) \in N_{i0}$ is determined by the equation $x_1 = it_1$.

Remark 4.12. If $\text{Exp}(\lambda, t_1) \in S_6$ for some $(\lambda, t) \in N$, then $\varepsilon^6(\lambda) = \lambda$.

Proof. It follows from Lemmas 4.8, 4.9, 4.10 that the inclusion $\text{Exp}(\lambda, t_1) \in S_6$ is possible only in the following two cases:

- (1) $\lambda \in C_6$, $\tau = \theta + ct/2 = \pm\pi/2$,
- (2) $\lambda \in C_7$, $\theta = \pm\pi/2$.

In both these cases we have $\varepsilon^6(\lambda) = \lambda$, since $\varepsilon^6: (\theta, c, \alpha) \mapsto (\pi - \theta_t, c_t, -\alpha) = (\pi - \theta_t, c, \alpha)$ and $\pi - \theta_t = \theta$. \square

5. CONCLUSION

We get a description of the global structure of the exponential mapping in the left-invariant sub-Riemannian problem on the Engel group. It was proved that restriction of this mapping to subdomains in the preimage and image of the exponential mapping cut out by the Maxwell strata corresponding to reflections is a diffeomorphism. Thus we reduced the problem to solving a system of algebraic equations. For any terminal point $q_1 = (x_1, y_1, z_1, v_1)$ with $x_1 \neq 0$ and $z_1 \neq 0$ there exists a unique optimal trajectory. Moreover it was proved that the cut time is equal to the first Maxwell time corresponding to reflections.

The cut locus in the sub-Riemannian problem on the Engel group will be described in a forthcoming article. We also plan to study sub-Riemannian spheres and their singularities. Developing software for computation of optimal solutions will allow us to solve the motion planning problem for generic control systems with 4 states and 2 linear inputs via nilpotent approximation (in particular, for the kinematic model of a car with trailer).

REFERENCES

- [1] Ardentov A.A., Sachkov Yu.L., Extremal trajectories in nilpotent sub-Riemannian problem on the Engel group, *Sbornik: Mathematics*, 2011, 202 (11), 1593-1615.
- [2] Ardentov A.A., Sachkov Yu.L., Conjugate points in nilpotent sub-Riemannian problem on the Engel group, *Journal of Mathematical Sciences*, 2013, 195 (3), 369-390.
- [3] Jurdjevic V., *Geometric Control Theory*, Cambridge University Press; 1997.
- [4] Agrachev A.A., Sachkov Yu.L., *Control Theory from the Geometric Viewpoint*, Vol. 87 of *Encyclopaedia of Mathematical Sciences*, Springer-Verlag, 2004.
- [5] Agrachev A.A., Barilari D., Boscain U., Introduction to Riemannian and sub-Riemannian geometry, Lecture notes, Preprint, 2014. <https://www.imj-prg.fr/~simon.davide.barilari/Notes.php>
- [6] Agrachev A.A., Bonnard B., Chyba M., Kupka I., Sub-Riemannian sphere in Martinet flat case, *J. ESAIM: Control, Optimisation and Calculus of Variations*, 1997, 2, 377-448.
- [7] Sachkov Yu.L., Exponential map in the generalized Dido problem, *Sbornik: Mathematics*, 2003, 194 (9), 1331-1359.
- [8] Sachkov Yu.L., Discrete symmetries in the generalized Dido problem, *Sbornik: Mathematics*, 2006, 197 (2), 235-257
- [9] Sachkov Yu.L., The Maxwell set in the generalized Dido problem, *Sbornik: Mathematics*, 2006, 197 (4), 595-621
- [10] Sachkov Yu.L., Complete description of the Maxwell strata in the generalized Dido problem, *Sbornik: Mathematics*, 2006, 197(6), 901-950
- [11] Moiseev I., Sachkov Yu.L., Maxwell strata in sub-Riemannian problem on the group of motions of a plane, *ESAIM: COCV*, 2010, 16, 380-399.
- [12] Sachkov Yu.L., Conjugate and cut time in sub-Riemannian problem on the group of motions of a plane, *ESAIM: COCV*, 2010, 16, 1018-1039.
- [13] Sachkov Yu.L., Cut locus and optimal synthesis in the sub-Riemannian problem on the group of motions of a plane, *ESAIM: Control, Optimisation and Calculus of Variations*, 2011, 17, 293-321.
- [14] Sachkov Yu.L., Maxwell strata in Euler's elastic problem, *Journal of Dynamical and Control Systems*, 2008, 14(2), 169-234.

- [15] Sachkov Yu.L., Conjugate points in Euler's elastic problem, *Journal of Dynamical and Control Systems*, Springer, New York, 2008, 14(3), 409-439.
- [16] Sachkov Yu.L., Sachkova E.F., Exponential mapping in Euler's elastic problem, *Journal of Dynamical and Control Systems*, Vol. 20 (2014), in print.
- [17] Agrachev A.A., Geometry of optimal control problems and Hamiltonian systems. In: *Nonlinear and Optimal Control Theory*, Lecture Notes in Mathematics. CIME, 1932, Springer Verlag, 2008, 1-59.
- [18] Krantz S. G., Parks H. R., *The Implicit Function Theorem: History, Theory, and Applications*, Birkauer, 2001.
- [19] Boscain U., Rossi F., Invariant Carnot-Caratheodory metrics on S^3 , $SO(3)$, $SL(2)$ and Lens Spaces. *SIAM, Journal on Control and Optimization*, 2008, 47, 1851-1878.
- [20] Butt Y.A., Sachkov Yu.L., Bhatti A.I., Parametrization of Extremal Trajectories in Sub-Riemannian Problem on Group of Motions of Pseudo Euclidean Plane, *Journal of Dynamical and Control Systems*, Vol. 20 (2014), No. 3 (July), 341-364.
- [21] Butt Y.A., Sachkov Yu.L., Bhatti A.I., Maxwell Strata and Conjugate Points in the Sub-Riemannian Problem on the Lie Group $SH(2)$, *submitted*.
- [22] Agrachev A.A., Barilari D.: Sub-Riemannian structures on 3D Lie groups, *Journal of Dynamical and Control Systems*, 2012, 18 (1), 21-44.
- [23] Almeida D.M., Sub-Riemannian Homogeneous Spaces of Engel Type, *Journal of Dynamical and Control Systems*, 2014, 20 (2), 149-166.
- [24] Vershik A.M., Gershkovich V.Ya., Nonholonomic dynamical systems, geometry of distributions and variational problems, *Dynamical Systems VII, Encyclopaedia of Mathematical Sciences (1990)*, 4-79
- [25] Montgomery R., *A tour of subriemannian geometries, their geodesics and applications*, AMS, 2002.
- [26] Pontryagin L.S., Boltayanskii V.G., Gamkrelidze R.V., Mishchenko E.F., *The mathematical theory of optimal processes*, John Wiley, New York/London, 1962.
- [27] Whittaker E.T., Watson G.N., *A Course of Modern Analysis*, Cambridge University Press, 1927.
- [28] Euler L., *Methodus inveniendi lineas curvas maximi minimive proprietate gaudentes, sive Solutio problematis isoperimetrici latissimo sensu accepti*, Lausanne, Geneva, 1744.
- [29] Love A.E.H., *A Treatise on the Mathematical Theory of Elasticity*, New York: Dover, 1927.