

Optimal control problem (*Lecture 4*)

Yuri Sachkov

Program Systems Institute
Russian Academy of Sciences
Pereslavl-Zalessky, Russia
yusachkov@gmail.com

«Geometric control theory, sub-Riemannian geometry, and their applications»

Mini-course in Fudan University, Shanghai, China

25 March 2022

3. *Seeing the Ox:*

On a yonder branch perches a nightingale cheerfully singing;
The sun is warm, and a soothing breeze blows, on the bank the willows are
green;
The ox is there all by himself, nowhere is he to hide himself;
The splendid head decorated with stately horns what painter can reproduce
him?

Pu-ming, "The Ten Oxherding Pictures"



Reminder: Plan of the previous lecture

1. Proof of the Orbit theorem.
2. Corollaries of the Orbit theorem:
 - Rashevskii–Chow theorem,
 - Lie algebra rank condition,
 - Frobenius theorem.
3. Krener's theorem.

Plan of this lecture

1. Statement of optimal control problem
2. Existence of optimal controls
3. Pontryagin maximum principle
4. Solution to examples of optimal control problems
5. Sub-Riemannian problems

Statement of optimal control problem

- We consider the following *optimal control problem*:

$$\dot{q} = f(q, u), \quad q \in M, \quad u \in U \subset \mathbb{R}^m, \quad (1)$$

$$q(0) = q_0, \quad q(t_1) = q_1, \quad (2)$$

$$J[u] = \int_0^{t_1} \varphi(q, u) dt \rightarrow \min, \quad (3)$$

t_1 fixed or free.

- A solution $q(t)$, $t \in [0, t_1]$, to this problem is said to be *(globally) optimal*.
- The following assumptions are made for the dynamics $f(q, u)$:
 - the mapping $q \mapsto f(q, u)$ is smooth for any $u \in U$,
 - the mapping $(q, u) \mapsto f(q, u)$ is continuous for any $q \in M$, $u \in \text{cl}(U)$,
 - the mapping $(q, u) \mapsto \frac{\partial f}{\partial q}(q, u)$ is continuous for any $q \in M$, $u \in \text{cl}(U)$.
- The same assumptions are made for the function $\varphi(q, u)$ that determines the cost functional J .
- Admissible control is $u \in L^\infty([0, t_1], U)$.

Reduction to the study of attainable sets

- In order to include the functional J into dynamics of the system, introduce a new variable equal to the running value of the cost functional along a trajectory $q_u(t)$:

$$y(t) = \int_0^t \varphi(q, u) dt.$$

- Respectively, we introduce an extended state $\hat{q} = \begin{pmatrix} y \\ q \end{pmatrix} \in \mathbb{R} \times M$ that satisfies an *extended control system*

$$\frac{d\hat{q}}{dt} = \begin{pmatrix} \dot{y} \\ \dot{q} \end{pmatrix} = \begin{pmatrix} \varphi(q, u) \\ f(q, u) \end{pmatrix} =: \hat{f}(\hat{q}, u).$$


- The boundary conditions for this system are

$$\hat{q}(0) = \begin{pmatrix} 0 \\ q_0 \end{pmatrix}, \quad \hat{q}(t_1) = \begin{pmatrix} J \\ q_1 \end{pmatrix}.$$

Reduction to the study of attainable sets

- A trajectory $q_{\bar{u}}(t)$ is optimal for the optimal control problem with fixed time t_1 if and only if the corresponding trajectory $\hat{q}_{\bar{u}}(t)$ of the extended system comes to a point (y_1, q_1) of the attainable set $\hat{\mathcal{A}}_{(0, q_0)}(t_1)$ such that

$$\hat{\mathcal{A}}_{(0, q_0)}(t_1) \cap \{(y, q_1) \mid y < y_1\} = \emptyset.$$



- For the problem with free terminal time an analogous condition is written for the attainable set $\hat{\mathcal{A}}_{(0, q_0)}$.

Filippov's theorem

Corollary

If the attainable set $\widehat{\mathcal{A}}_{(0,q_0)}(t_1)$ is compact and $q_1 \in \mathcal{A}_{q_0}(t_1)$, then the optimal control problem (1)–(3) with fixed time t_1 has a solution.

Theorem (Filippov)

Suppose that control system (1) satisfies the hypotheses:

- (1) the set U is compact*
- (2) the set $f(q, U)$ is convex for all $q \in M$*
- (3) there exists a compact set $K \subset M$ such that for all $q \in M \setminus K$, $u \in U$ there holds the equality $f(q, u) = 0$.*

Then the attainable sets $\mathcal{A}_{q_0}(t)$, $\mathcal{A}_{q_0}(\leq t)$ are compact for any $q_0 \in M$, $t > 0$.

Existence of optimal controls in optimal control problem

Corollary

Let the optimal control problem (1)–(3) satisfy the hypotheses:

- (1) the set U is compact
- (2) the set $\left\{ \begin{pmatrix} \varphi(q, u) \\ f(q, u) \end{pmatrix} \mid u \in U \right\}$ is convex for all $q \in M$
- (3) there exists a compact set $K \subset \mathbb{R} \times M$ such that $\hat{\mathcal{A}}_{(0, q_0)}(t_1) \subset K$
- (4) $q_1 \in \mathcal{A}_{q_0}(t_1)$.

Then the problem (1)–(3) with fixed time t_1 has a solution.

Proof of the existence conditions for optimal control problem

- *Proof.* There exists a compact set $K' \subset \mathbb{R} \times M$ such that $K \subset \text{int } K'$. Take a function $a \in C^\infty(\mathbb{R} \times M)$ such that

$$a|_K \equiv 1, \quad a|_{(\mathbb{R} \times M) \setminus K'} \equiv 0.$$

- Consider a new extended control system:

$$\frac{d\hat{q}}{dt} = a(\hat{q})f(\hat{q}, u), \quad \hat{q} \in \mathbb{R} \times M, \quad u \in U.$$

- This system has compact attainable sets for time t_1 , which coincide with the corresponding attainable sets of the extended system.
- Then optimal control problem (1)–(3) has a solution (by Filippov's theorem). □

Existence of solutions to time-optimal problem

Now consider a *time-optimal problem*

$$\dot{q} = f(q, u), \quad q \in M, \quad u \in U \subset \mathbb{R}^m, \quad (4)$$

$$q(0) = q_0, \quad q(t_1) = q_1, \quad (5)$$

$$t_1 \rightarrow \min. \quad (6)$$

Corollary

Let the following conditions hold:

- (1) the set U is compact
- (2) the set $f(q, U)$ is convex for all $q \in M$
- (3) there exist $t_1 > 0$ and a compact set $K \subset M$ such that

$$q_1 \in \mathcal{A}_{q_0}(\leq t_1) \subset K.$$

Then time-optimal problem (4)–(6) has a solution.

Elements of symplectic geometry

- Let M be an n -dimensional smooth manifold. Then the disjoint union of its tangent spaces $TM = \bigsqcup_{q \in M} T_q M = \{(q, v) \mid q \in M, v \in T_q M\}$ is called its *tangent bundle*.
- If (q_1, \dots, q_n) are local coordinates on M , then any tangent vector $v \in T_q M$ has a decomposition $v = \sum_{i=1}^n v_i \frac{\partial}{\partial q_i}$. So $(q_1, \dots, q_n; v_1, \dots, v_n)$ are local coordinates on TM , which is thus a $2n$ -dimensional smooth manifold.
- For any point $q \in M$, the dual space $(T_q M)^* = T_q^* M$ is called the *cotangent space* to M at q . Thus $T_q^* M$ consists of linear forms on $T_q M$. The disjoint union $T^* M = \bigsqcup_{q \in M} T_q^* M = \{(q, p) \mid q \in M, p \in T_q^* M\}$ is called the *cotangent bundle*.
- If (q_1, \dots, q_n) are local coordinates on M , then any covector $\lambda \in T^* M$ has a decomposition $\lambda = \sum_{i=1}^n p_i dq_i$. Thus $(q_1, \dots, q_n; p_1, \dots, p_n)$ are local coordinates on $T^* M$ called the *canonical coordinates*. So $T^* M$ is a smooth $2n$ -dimensional manifold.
- The *canonical projection* is the mapping $\pi: T^* M \rightarrow M, \quad T_q^* M \ni \lambda \mapsto q \in M$.

Elements of symplectic geometry

- The *Liouville (tautological) differential 1-form* $s \in \Lambda^1(T^*M)$ is defined as follows:

$$\langle s_\lambda, w \rangle = \langle \lambda, \pi_* w \rangle, \quad \lambda \in T^*M, \quad w \in T_\lambda(T^*M).$$

In the canonical coordinates on T^*M , $s = p dq$.

- The canonical *symplectic structure* on T^*M is the differential 2-form $\sigma = ds \in \Lambda^2(T^*M)$. In the canonical coordinates $\sigma = dp \wedge dq = \sum_{i=1}^n dp_i \wedge dq_i$.
- A *Hamiltonian (Hamiltonian function)* is an arbitrary function $h \in C^\infty(T^*M)$.
- The *Hamiltonian vector field* $\vec{h} \in \text{Vec}(T^*M)$ with the Hamiltonian function h is defined by the equality $dh = \sigma(\cdot, \vec{h})$. In the canonical coordinates:

$$h = h(q, p),$$

$$\vec{h} = \frac{\partial h}{\partial p} \frac{\partial}{\partial q} - \frac{\partial h}{\partial q} \frac{\partial}{\partial p} = \sum_{i=1}^n \left(\frac{\partial h}{\partial p_i} \frac{\partial}{\partial q_i} - \frac{\partial h}{\partial q_i} \frac{\partial}{\partial p_i} \right).$$

Elements of symplectic geometry

- The corresponding *Hamiltonian system of ODEs* is

$$\dot{\lambda} = \vec{h}(\lambda), \quad \lambda \in T^*M.$$

- In the canonical coordinates:

$$\begin{cases} \dot{q} = \frac{\partial h}{\partial p}, \\ \dot{p} = -\frac{\partial h}{\partial q}, \end{cases} \quad \text{or} \quad \begin{cases} \dot{q}_i = \frac{\partial h}{\partial p_i}, \\ \dot{p}_i = -\frac{\partial h}{\partial q_i}, \end{cases} \quad i = 1, \dots, n.$$

- The *Poisson bracket* of Hamiltonians $h, g \in C^\infty(T^*M)$ is the Hamiltonian $\{h, g\} \in C^\infty(T^*M)$ defined by the equalities

$$\{h, g\} = \vec{h}g = \sigma(\vec{h}, \vec{g}).$$

- In the canonical coordinates:

$$\{h, g\} = \frac{\partial h}{\partial p} \frac{\partial g}{\partial q} - \frac{\partial h}{\partial q} \frac{\partial g}{\partial p} = \sum_{i=1}^n \left(\frac{\partial h}{\partial p_i} \frac{\partial g}{\partial q_i} - \frac{\partial h}{\partial q_i} \frac{\partial g}{\partial p_i} \right).$$

Elements of symplectic geometry

Lemma

Let $a, b, c \in C^\infty(T^*M)$ and $\alpha, \beta \in \mathbb{R}$. Then:

- (1) $\{a, b\} = -\{b, a\}$,
- (2) $\{a, a\} = 0$,
- (3) $\{\{a, b\}, c\} + \{\{b, c\}, a\} + \{\{c, a\}, b\} = 0$,
- (4) $\{\alpha a + \beta b, c\} = \alpha\{a, c\} + \beta\{b, c\}$,
- (5) $\{ab, c\} = \{a, c\}b + a\{b, c\}$,
- (6) $[\vec{a}, \vec{b}] = \vec{d}$, $d = \{a, b\}$.

Theorem (Noether)

Let $a, h \in C^\infty(T^*M)$. Then

$$a(e^{t\vec{h}}(\lambda)) \equiv \text{const} \quad \Leftrightarrow \quad \{h, a\} = 0.$$

Elements of symplectic geometry

Now we describe the last construction of symplectic geometry necessary for us — *linear on fibers of T^*M Hamiltonians*. Let $X \in \text{Vec}(M)$. The corresponding linear on fibers of T^*M Hamiltonian is defined as follows: $h_X(\lambda) = \langle \lambda, X(q) \rangle$, $q = \pi(\lambda)$.

In the canonical coordinates:

$$X = \sum_{i=1}^n X_i \frac{\partial}{\partial q_i}, \quad h_X(q, p) = \sum_{i=1}^n p_i X_i.$$

Lemma

Let $X, Y \in \text{Vec}(M)$. Then:

- (1) $\{h_X, h_Y\} = h_{[X, Y]}$,
- (2) $[\vec{h}_X, \vec{h}_Y] = \vec{h}_{[X, Y]}$,
- (3) $\pi_* \vec{h}_X = X$.

The vector field $\vec{h}_X \in \text{Vec}(T^*M)$ is called the *Hamiltonian lift* of the vector field $X \in \text{Vec}(M)$.

Hamiltonians of Pontryagin maximum principle

- Return to the optimal control problem

$$\dot{q} = f(q, u), \quad q \in M, \quad u \in U \subset \mathbb{R}^m,$$

$$q(0) = q_0, \quad q(t_1) = q_1,$$

$$J = \int_0^{t_1} \varphi(q, u) dt \rightarrow \min,$$

t_1 fixed.

- Define a family of *Hamiltonians of PMP*

$$h_u^\nu(\lambda) = \langle \lambda, f(q, u) \rangle + \nu \varphi(q, u), \quad \nu \in \mathbb{R}, \quad u \in U, \quad \lambda \in T^*M, \quad q = \pi(\lambda).$$

Statement of Pontryagin maximum principle

Theorem (PMP)

If a control $u(t)$ and the corresponding trajectory $q(t)$, $t \in [0, t_1]$, are optimal, then there exist a curve $\lambda_t \in \text{Lip}([0, t_1], T^*M)$, $\lambda_t \in T_{q(t)}^*M$, and a number $\nu \leq 0$ such that the following conditions hold for almost all $t \in [0, t_1]$:

- (1) $\dot{\lambda}_t = \vec{h}_{u(t)}^\nu(\lambda_t)$,
- (2) $h_{u(t)}^\nu(\lambda_t) = \max_{w \in U} h_w^\nu(\lambda_t)$,
- (3) $(\lambda_t, \nu) \neq (0, 0)$.

If the terminal time t_1 is free, then the following condition is added to (1)–(3):

- (4) $h_{u(t)}^\nu(\lambda_t) \equiv 0$.

A curve λ_t that satisfies PMP is called an *extremal*, a curve $q(t)$ — an *extremal trajectory*, a control $u(t)$ — an *extremal control*.

Time-optimal problem

- Let us apply PMP to the *time-optimal problem*

$$\dot{q} = f(q, u), \quad q \in M, \quad u \in U,$$

$$q(0) = q_0, \quad q(t_1) = q_1,$$

$$t_1 = \int_0^{t_1} 1 dt \rightarrow \min.$$

- The Hamiltonian of PMP has the form $h_u^\nu(\lambda) = \langle \lambda, f(q, u) \rangle + \nu$. Introduce the *shortened Hamiltonian* $g_u(\lambda) = \langle \lambda, f(q, u) \rangle$.
- Then the statement of PMP for the time-optimal problem takes the form:
 - $\dot{\lambda}_t = \vec{h}_{u(t)}^\nu(\lambda_t) = \vec{g}_{u(t)}(\lambda_t),$
 - $h_{u(t)}^\nu(\lambda_t) = \max_{w \in U} h_w^\nu(\lambda_t) \Leftrightarrow g_{u(t)}(\lambda_t) = \max_{w \in U} g_w(\lambda_t),$
 - $\lambda_t \neq 0,$
 - $h_{u(t)}^\nu(\lambda_t) \equiv 0 \Leftrightarrow g_{u(t)}(\lambda_t) \equiv \text{const} \geq 0.$

The case of smooth maximized Hamiltonian

Denote the *maximized normal Hamiltonian of PMP*

$$H(\lambda) = \max_{u \in U} h_u^{-1}(\lambda), \quad \lambda \in T^*M.$$

Theorem

Let $H \in C^2(T^*M)$. Then a curve λ_t is a normal extremal iff it is a trajectory of the Hamiltonian system $\dot{\lambda}_t = \vec{H}(\lambda_t)$.

Example: Stopping a train

- We have the time-optimal problem

$$\begin{aligned}\dot{x}_1 &= x_2, & \dot{x}_2 &= u, & x &= (x_1, x_2) \in \mathbb{R}^2, & |u| &\leq 1, \\ x(0) &= x^0, & x(t_1) &= x^1 = (0, 0), & t_1 &\rightarrow \min.\end{aligned}$$

- The right-hand side of the control system $f(x, u) = (x_2, u)$ satisfies the bound

$$|f(x, u)| = \sqrt{x_2^2 + u^2} \leq \sqrt{x_2^2 + 1} \leq |x| + 1,$$

thus $r = x^2$ satisfies the differential inequality

$\dot{r} = 2\langle x, \dot{x} \rangle = 2\langle x, f(x, u) \rangle \leq 2(r + 1)$. So $r(t) \leq e^{2t}(r_0 + 1)$, thus attainable sets satisfy the a priori bound

$$\mathcal{A}_{x^0}(\leq t) \subset \left\{ x \in \mathbb{R}^2 \mid |x| \leq e^t \sqrt{(x^0)^2 + 1} \right\}.$$

- Therefore we can assume that there exists a compact set $K \subset \mathbb{R}^2$ such that the right-hand side of the control system vanishes outside of K (one of conditions of the Filippov theorem).

Example: Stopping a train

- As we showed, $x^1 = (0, 0) \in \mathcal{A}_{x^0}$ for any $x^0 \in \mathbb{R}^2$.
- The set of control parameters U is compact, and the set of admissible velocity vectors $f(x, U)$ is convex for any $x \in \mathbb{R}^2$. All hypotheses of the Filippov theorem are satisfied, thus optimal control exists.
- We apply PMP using the canonical coordinates (p_1, p_2, x_1, x_2) on $T^*\mathbb{R}^2$. We decompose a covector $\lambda = p_1 dx_1 + p_2 dx_2 \in T^*\mathbb{R}^2$, then the shortened Hamiltonian of PMP reads $h_u(\lambda) = p_1 x_2 + p_2 u$, and the Hamiltonian system $\dot{\lambda} = \vec{h}_u(\lambda)$ reads

$$\begin{aligned}\dot{x}_1 &= x_2, & \dot{p}_1 &= 0, \\ \dot{x}_2 &= u, & \dot{p}_2 &= -p_1.\end{aligned}$$

- The maximality condition of PMP has the form

$$h_u(\lambda) = p_1 x_2 + p_2 u \rightarrow \max_{|u| \leq 1}$$

and the nontriviality condition is $(p_1(t), p_2(t)) \neq (0, 0)$.

Example: Stopping a train

- The maximality condition yields:

$$p_2(t) > 0 \Rightarrow u(t) = 1, \quad p_2(t) < 0 \Rightarrow u(t) = -1.$$

- Thus extremal trajectories are the parabolas

$$x_1 = \pm \frac{x_2^2}{2} + C,$$

and the number of switchings (discontinuities) of control is not greater than 1.


- Let us construct such trajectories backward in time, starting from $x^1 = (0, 0)$:
 - the controls $u = 1$ and $u = -1$ generate two half-parabolas terminating at x^1 :

$$x_1 = \frac{x_2^2}{2}, \quad x_2 \leq 0 \quad \text{and} \quad x_1 = -\frac{x_2^2}{2}, \quad x_2 \geq 0,$$

- denote the union of these half-parabolas as Γ ,
- after one switching, parabolic arcs with $u = 1$ terminating at the half-parabola $x_1 = -\frac{x_2^2}{2}, \quad x_2 \geq 0$, fill the part of the plane \mathbb{R}^2 below the curve Γ ,
- similarly, after one switching, parabolic arcs with $u = -1$ fill the part of the plane over the curve Γ .

Example: Stopping a train

- So through each point of the plane \mathbb{R}^2 passes a unique extremal trajectory. In view of existence of optimal controls, the extremal trajectories are optimal.
- The optimal control found has explicit dependence on the current point of the plane: if $x_1 = \frac{x_2^2}{2}$, $x_2 \leq 0$, or if the point (x_1, x_2) lies below the curve Γ , then $u(x_1, x_2) = 1$, otherwise, $u(x_1, x_2) = -1$.



- Such a dependence $u(x)$ of optimal control on the current point x of the state space is called an *optimal synthesis*, it is the best possible form of solution to an optimal control problem.

Example: The Markov–Dubins car

- We have a time-optimal problem

$$\begin{aligned}\dot{x} &= \cos \theta, & q &= (x, y, \theta) \in \mathbb{R}_{x,y}^2 \times S_\theta^1 = M, \\ \dot{y} &= \sin \theta, & |u| &\leq 1, \\ \dot{\theta} &= u, \\ q(0) &= q_0 = (0, 0, 0), & q(t_1) &= q_1, \\ t_1 &\rightarrow \min.\end{aligned}$$

- The system is completely controllable.
- All conditions of the Filippov theorem are satisfied: U is compact, $f(q, U)$ are convex, the bound $|f(q, u)| \leq 2$ implies a priori bound of the attainable set. Thus optimal control exists.
- We apply PMP.

Example: The Markov–Dubins car

- The vector fields

$$f_0 = \cos \theta \frac{\partial}{\partial x} + \sin \theta \frac{\partial}{\partial y},$$

$$f_1 = \frac{\partial}{\partial \theta},$$

$$f_2 = [f_0, f_1] = \sin \theta \frac{\partial}{\partial x} - \cos \theta \frac{\partial}{\partial y}$$

form a frame in $T_q M$.

- Define the corresponding linear on fibers of T^*M Hamiltonians:

$$h_i(\lambda) = \langle \lambda, f_i \rangle, \quad i = 0, 1, 2.$$

- The shortened Hamiltonian of PMP is

$$h_u(\lambda) = \langle \lambda, f_0 + u f_1 \rangle = h_0 + u h_1.$$

Example: The Markov–Dubins car

- The functions h_0, h_1, h_2 form a coordinate system on T_q^*M , and we write the Hamiltonian system of PMP in the non-canonical parametrization (h_0, h_1, h_2, q) of T^*M :

$$\dot{h}_0 = \vec{h}_u h_0 = \{h_0 + uh_1, h_0\} = -uh_2, \quad (7)$$

$$\dot{h}_1 = \{h_0 + uh_1, h_1\} = h_2, \quad (8)$$

$$\dot{h}_2 = \{h_0 + uh_1, h_2\} = uh_0, \quad (9)$$

$$\dot{q} = f_0 + uf_1.$$


- The maximality condition $h_u(\lambda) = h_0 + uh_1 \rightarrow \max_{|u| \leq 1}$ implies that if $h_1(\lambda_t) \neq 0$, then $u(t) = \text{sgn } h_1(\lambda_t)$.
- Consider the case where the control is not determined by PMP: $h_1(\lambda_t) \equiv 0$ (this case is called *singular*). Then (8) gives $h_2(\lambda_t) \equiv 0$, thus $h_0(\lambda_t) \neq 0$ by the nontriviality condition of PMP, so $u(t) \equiv 0$ by (9). The corresponding extremal trajectory $(x(t), y(t))$ is a straight line.

Example: The Markov–Dubins car

- If $u(t) = \pm 1$, then the extremal trajectory $(x(t), y(t))$ is an arc of a unit circle.
- One can show that optimal trajectories have one of the following two types:
 1. arc of unit circle + line segment + arc of unit circle
 2. concatenation of three arcs of unit circles; in this case, if a, b, c are the times along the first, second, and third arc respectively, then $\pi < b < 2\pi$, $\min\{a, c\} < b$, and $\max\{a, c\} < b$.
- If boundary conditions are far one from another, then the optimal trajectory has type 1 and can explicitly be constructed as follows. Draw two unit circles that satisfy the initial condition and two unit circles that satisfy the terminal condition. Draw four common tangents to the initial circles and the terminal circles, with account of direction of motion along the circles determined by the boundary conditions. Among the four constructed extremal trajectories, find the shortest one. It is the optimal trajectory.
- The optimal synthesis for the Markov–Dubins car is known, but it is rather complicated.

Example: Control of linear oscillator

- Optimal trajectories are concatenations of circular arcs.
- The optimal synthesis:



Sub-Riemannian structures and minimizers

- A *sub-Riemannian structure* on a smooth manifold M is a pair (Δ, g) , where

$$\Delta = \{\Delta_q \subset T_q M \mid q \in M\}$$

is a distribution on M and

$$g = \{g_q \text{ inner product in } \Delta_q \mid q \in M\}$$

is an *inner product* (nondegenerate positive definite quadratic form) on Δ .

- The spaces Δ_q and inner products g_q depend smoothly on $q \in M$, and $\dim \Delta_q \equiv \text{const}$.
- A curve $q \in \text{Lip}([0, t_1], M)$ is called *horizontal (admissible)* if

$$\dot{q}(t) \in \Delta_{q(t)} \text{ for almost all } t \in [0, t_1].$$

- The *sub-Riemannian length* of a horizontal curve $q(\cdot)$ is defined as

$$l(q(\cdot)) = \int_0^{t_1} \sqrt{g(\dot{q}, \dot{q})} dt.$$

Sub-Riemannian structures and minimizers

- The *sub-Riemannian (Carnot–Carathéodory) distance* between points $q_0, q_1 \in M$ is

$$d(q_0, q_1) = \inf\{l(q(\cdot)) \mid q(\cdot) \text{ horizontal, } q(0) = q_0, q(t_1) = q_1\}.$$

- A horizontal curve $q(\cdot)$ is called a *sub-Riemannian length minimizer* if

$$l(q(\cdot)) = d(q(0), q(t_1)).$$

- Thus length minimizers are solutions to a *sub-Riemannian optimal control problem*:

$$\dot{q}(t) \in \Delta_{q(t)},$$

$$q(0) = q_0, \quad q(t_1) = q_1,$$

$$l(q(\cdot)) \rightarrow \min.$$

- Suppose that a sub-Riemannian structure (Δ, g) has a *global orthonormal frame* $f_1, \dots, f_k \in \text{Vec}(M)$:

$$\Delta_q = \text{span}(f_1(q), \dots, f_k(q)), \quad q \in M, \quad g(f_i, f_j) = \delta_{ij}, \quad i, j = 1, \dots, k.$$

Sub-Riemannian structures and minimizers

- Then the optimal control problem for sub-Riemannian minimizers takes the standard form:

$$\dot{q} = \sum_{i=1}^k u_i f_i(q), \quad q \in M, \quad u = (u_1, \dots, u_k) \in \mathbb{R}^k, \quad (10)$$

$$q(0) = q_0, \quad q(t_1) = q_1, \quad (11)$$

$$l = \int_0^{t_1} \left(\sum_{i=1}^k u_i^2 \right)^{1/2} dt \rightarrow \min. \quad (12)$$

- The sub-Riemannian length does not depend on parametrization of a horizontal curve $q(t)$. Namely, if

$$\tilde{q}(s) = q(t(s)), \quad t(\cdot) \in \text{Lip}([0, s_1], [0, t_1]), \quad t'(s) > 0,$$

is a reparametrization of a curve $q(t)$, then $l(\tilde{q}(\cdot)) = l(q(\cdot))$.

Sub-Riemannian structures and minimizers

- Along with the length functional, it is convenient to consider the *energy* functional

$$J(q(\cdot)) = \frac{1}{2} \int_0^{t_1} g(\dot{q}, \dot{q}) dt.$$

- Denote $\|\dot{q}\| = \sqrt{g(\dot{q}, \dot{q})}$.

Sub-Riemannian structures and minimizers

Lemma

Let the terminal time t_1 be fixed. Then minimizers of energy are exactly length minimizers of constant velocity:

$$J(q(\cdot)) \rightarrow \min \quad \Leftrightarrow \quad l(q(\cdot)) \rightarrow \min, \quad \|\dot{q}\| = \text{const}.$$

Proof.

By the Cauchy–Schwarz inequality,

$$(l(q(\cdot)))^2 = \left(\int_0^{t_1} \|\dot{q}\| \cdot 1 \, dt \right)^2 \leq \int_0^{t_1} \|\dot{q}\|^2 \, dt \cdot \int_0^{t_1} 1^2 \, dt = 2J(q(\cdot)) t_1,$$

moreover, equality is attained here only for $\|\dot{q}\| = \text{const}$.

It is obvious that on constant velocity curves the problems $l \rightarrow \min$ and $J \rightarrow \min$ are equivalent. And for $\|\dot{q}\| \neq \text{const}$ we have $l < 2t_1 J$, i.e., J does not attain minimum. \square